

# Papéis Avulsos de Zoologia

Museu de Zoologia da Universidade de São Paulo

Volume 50(1):1-29, 2010

[www.mz.usp.br/publicacoes](http://www.mz.usp.br/publicacoes)

[www.revistasusp.sibi.usp.br](http://www.revistasusp.sibi.usp.br)

[www.scielo.br/paz](http://www.scielo.br/paz)

ISSN impresso: 0031-1049

ISSN on-line: 1807-0205

## PATTERNS OF GEOGRAPHIC DISTRIBUTION AND CONSERVATION OF THE OPEN-HABITAT AVIFAUNA OF SOUTHEASTERN BRAZILIAN MOUNTAINTOPS (CAMPOS RUPESTRES AND CAMPOS DE ALTITUDE)

MARCELO FERREIRA DE VASCONCELOS<sup>1,2</sup>

MARCOS RODRIGUES<sup>1,3</sup>

### ABSTRACT

*Montane open-habitats of southeastern Brazil are represented by the campos rupestres (principally in the Espinhaço Range) and by the campos de altitude (in the Serra do Mar and Serra da Mantiqueira). In spite of the occurrence of endemic species in both vegetation types, an analysis and synthesis of their bird communities have never been conducted. In this paper, we present an avifaunal survey of these areas, describe patterns of geographic distribution, and comment on the conservation of those open-habitats and their avifauna. A total of 231 bird species was recorded in the open-habitats of southeastern Brazilian mountaintops. In the campos rupestres, 205 species were recorded, while in the campos de altitude, the total was 123 species. Five patterns of distribution are recognizable among birds occurring in these habitats: non-endemic (191 species), Atlantic Forest endemics (26 species), Cerrado endemics (6 species), Caatinga endemic (1 species), and montane open-habitat endemics (7 species). In spite of the presence of several protected areas in those regions, the existing reserves do not guarantee the conservation of their important vegetation types and their avifaunas under current low levels of implementation. Since several endemic and threatened bird species live in the campos rupestres and campos de altitude, more efforts must be directed for their conservation.*

KEYWORDS: Avifauna; *Campos rupestres*; *Campos de altitude*; Conservation; Distribution.

### INTRODUCTION

Montane open-habitats in southeastern Brazil are recognized as important centres of endemism for Neotropical flora and fauna (Eiten, 1992; Silva, 1995, 1997, 1998; Giulietti *et al.*, 1997; Sick, 1997;

Stattersfield *et al.*, 1998; Safford, 1999a; Silva & Bates, 2002; Gonçalves *et al.*, 2007). Those habitats generally occur at higher elevations than the Cerrado and the Caatinga in their northern and western distribution and above the tree line in the Atlantic Forest region to the south and east.

1. Departamento de Zoologia, ICB, Universidade Federal de Minas Gerais, Caixa Postal 486, 30123-970, Belo Horizonte, MG, Brasil.

2. Corresponding author: Rua Paraíba, 740, Apartamento 501, 30130-140, Belo Horizonte, MG, Brasil. Fone: +55 31 3425-4673.  
E-mail: [mfvasconcelos@gmail.com](mailto:mfvasconcelos@gmail.com).

3. E-mail: [ornito@icb.ufmg.br](mailto:ornito@icb.ufmg.br).

Montane open-habitats include the *campos rupestres* scattered along the Espinhaço Range and the *campos de altitude* dotting the ridges of the eastern mountains of the Serra do Mar, Serra da Mantiqueira, and associated ranges (Martinelli & Orleans e Bragança, 1996; Giuliatti *et al.*, 1997; Safford, 1999a; Menezes & Giuliatti, 2000). Both habitats harbor a wide variety of physiognomies, ranging from open grasslands to habitats with a dense cover of scrub and small trees. Rock outcrops and marshy areas also occur in both formations, which have many typical and endemic plants, such as bromeliads (Bromeliaceae), “semprevivas” (Eriocaulaceae), “canelas-de-ema” (Velloziaceae), bamboos and grasses (Poaceae) (Andrade *et al.*, 1986; Eiten, 1992; Meguro *et al.*, 1994; Martinelli & Orleans e Bragança, 1996; Safford, 1999a; Menezes & Giuliatti, 2000; Conceição & Giuliatti, 2002; Benites *et al.*, 2003; Caiafa & Silva, 2005; Conceição & Pirani, 2005; Ribeiro *et al.*, 2007).

Although *campos rupestres* and *campos de altitude* can present similar landscapes and share some common plant genera and species, they show important differences in the biogeographic affinities of their floras (Menezes & Giuliatti, 1986; Eiten, 1992; Harley, 1995; Martinelli & Orleans e Bragança, 1996; Giuliatti *et al.*, 1997; Rizzini, 1997; Safford, 1999a, 2007). The *campos rupestres* flora has affinities with those of central Brazilian Plateau, the Guianas Shield, and the restingas along the Atlantic Ocean coast of South America (Steyermark, 1986; Giuliatti & Pirani, 1988; Giuliatti & Hensold, 1990; Harley, 1995; Menezes & Giuliatti, 2000; Alves, R.J.V. *et al.*, 2007), but with a high number of endemic Bromeliaceae, Eriocaulaceae, Orchidaceae, and Velloziaceae (Eiten, 1992; Menezes & Giuliatti, 2000; Versieux & Wendt, 2007).

On the other hand, *campos de altitude* show a remarkable affinity with the Andean region and also with southern Brazilian grasslands (Harley, 1995; Martinelli & Orleans e Bragança, 1996; Safford, 1999a, 2007). In fact, *campos de altitude* have been considered as ‘Brazilian Páramos’ because their floristic, physiognomic, climatic, and edaphic features are so similar to those of the Andes and the high Central American mountains (Harley, 1995; Safford, 1999a, b, 2007).

Furthermore, biogeographical affinities have been recognized between the flora and fauna of southeastern Brazilian mountains and the Andean-Patagonian region (Sick, 1970, 1985, 1997; Simpson-Vuilleumier, 1971; Simpson, 1979; Vielliard, 1990a, 1996; Safford, 1999a, 2007).

It is assumed that Brazilian mountains were colonized by Andean elements during Pleistocene

glaciations, when climatic-vegetational connections between both regions were likely (Simpson-Vuilleumier, 1971; Simpson, 1979; Sick, 1970, 1985, 1997; Vielliard, 1990a). Those cooler periods would have led to the lowering of the tree line, increasing the latitudinal distribution of open-habitats in southeastern Brazil, similar to what is believed to have occurred in the Andean “páramos” (van der Hammen, 1974; Simpson-Vuilleumier, 1971; Simpson, 1979; Hooghiemstra & Cleef, 1995; Behling, 1998; Ledru *et al.*, 1998; Safford, 1999a). Also, during Late Glacial times, subtropical grasslands expanded from latitudes of about 28°/27°S to at least 20°S (Behling, 2002). During interglacial periods, those montane habitats would have retreated to cooler upland areas, explaining the isolated occurrence of some Andean-Patagonian taxa on southeastern Brazilian mountaintops (Simpson, 1979; Safford, 1999a). This model supports the notion that the high-altitude avifauna of southeastern Brazil has a biogeographical influence of Andean-Patagonian elements, with the putative dispersion of those taxa from the Andes and Patagonia to southeastern Brazil during Quaternary glacial periods (Sick, 1970, 1985, 1997; Vielliard, 1990a, 1996; Willis, 1992).

Despite the occurrence of some endemic birds in the montane open-habitats of southeastern Brazil (Sick, 1970, 1985, 1997; Silva, 1995; Melo-Júnior *et al.*, 1998; Stattersfield *et al.*, 1998; Vasconcelos *et al.*, 2003; Gonzaga *et al.*, 2007; Vasconcelos, 2008b), their biogeography is poorly known, with only sparse surveys and checklists (*e.g.* Miranda-Ribeiro, 1906, 1923; Lüderwaldt, 1909; Peixoto-Velho, 1923a, b; Holt, 1928; Pinto, 1951, 1954; Willis & Oniki, 1991; Parrini *et al.*, 1999; Simon *et al.*, 1999; Vasconcelos, 1999a, 2003, 2008a; Melo-Júnior *et al.*, 2001; Vasconcelos & Melo-Júnior, 2001). Bird communities from the *campos rupestres* and *campos de altitude* have never been subject to a thorough analysis and biogeographical synthesis. A good example of how poorly surveyed are these areas is the recent description of *Formicivora grantsaui* in the Chapada Diamantina (Gonzaga *et al.*, 2007), a relatively well-known region (Funch, 1999; Parrini *et al.*, 1999; Carvalhaes, 2001).

In his classic review of the physical environment and flora of the *campos de altitude*, Safford (1999a) recommended flora and fauna inventories as a first priority for research and conservation of this habitat. Following his lead, the goals of this paper are: 1) to review and present an ornithological synthesis of the *campos rupestres* and *campos de altitude*; 2) to describe patterns of geographic distribution of the montane open-habitat avifauna of southeastern Brazil; 3) to discuss the conservation of those habitats and their avifauna.

**MATERIAL AND METHODS**


We included in our survey those species that live in open natural habitats in the mountain areas of southeastern Brazil from Bahia south to São Paulo (Fig. 1). Those include marshes, grasslands, bamboo stands (*Chusquea* spp.), rocky outcrops covered with rupicolous herbs and shrubs, or with clumped shrubs and sparse low trees. Both *campos rupestres* and *campos de altitude* occur over a wide variety of rock substrates, such as granite, gneiss, quartzite, sandstone, and even iron ore. For physical and floristic characterizations of these habitats, see Andrade *et al.* (1986), Menezes & Giulietti (1986, 2000), Eiten (1992), Alves & Kolbek (1994), Meguro *et al.* (1994), Harley (1995), Martinnelli & Orleans e Bragança (1996), Rizzini (1997), Safford (1999a), Conceição & Giulietti (2002), Dias *et al.* (2002), Benites *et al.* (2003, 2007), Vincent (2004), Caiafa & Silva (2005, 2007), Conceição & Pirani (2005), Conceição *et al.* (2007), Jacobi *et al.* (2007), and Ribeiro *et al.* (2007).

In this survey, we did not include species living strictly in forested habitats associated with the *campos*

*rupestres* or the *campos de altitude*, such as gallery, cloud, and montane forests. Nevertheless, some typical forest birds do use dense clumps of shrubs or bamboos in those montane open-habitats (e.g. *Mackenziaena leachii*, *Drymophila genei*, *Hemitriccus obsoletus*, *Phylloscartes difficilis*, and *Haplospiza unicolor*), and they were included in this revision. We also excluded bird species that live only in the different physiognomies of the Cerrado, such as cerradão, cerrado “sensu stricto”, parque de cerrado, and campo sujo (see Ribeiro & Walter, 1998 for a detailed description of these habitat types).

We selected different areas of *campos rupestres* and *campos de altitude* for our analysis (Fig. 1); the Chapada Diamantina region (Dias Coelho) was considered our northernmost site and Serra da Mantiqueira (Atibaia) the southern limit of the study area (Fig. 1). Localities surveyed in this study are summarized in Table 1. They were separated by mountain range (Espinhaço Range, Serra da Mantiqueira, and Serra do Mar) and type of habitat (*campo rupestre* or *campo de altitude*). All localities in the Espinhaço Range are covered by *campos rupestres*, and all localities in the Serra do Mar are represented by *campos de altitude*. The majority of the localities in the Serra da Mantiqueira are covered by *campos de altitude*, except Serra de São José and Serra do Lenheiro, which are covered by *campos rupestres* (see Alves & Kolbelk, 1994, 2009).

Due to the scarcity of data on birds of Brazilian montane open-habitats, we used all information gathered during fieldwork between 1993 and 2008. Voucher specimens were collected with mist-nets and shotguns, and deposited in the Coleção Ornitológica do Departamento de Zoologia da Universidade Federal de Minas Gerais, Belo Horizonte (DZUFMG). We also checked published bird records and data from specimens deposited in the following museums and collections: American Museum of Natural History, New York (AMNH); Museu de Biologia Mello Leitão, Santa Teresa (MBML); Museu de Ciências Naturais da Pontifícia Universidade Católica de Minas Gerais, Belo Horizonte (MCN); Museu de Ciências e Tecnologia da Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre (MCP); Museu de História Natural de Taubaté, Taubaté (MHNT); Museu Paraense Emílio Goeldi, Belém (MPEG); Museu de Zoologia da Universidade de São Paulo, São Paulo (MZUSP); and Coleção Rolf Grantsau, São Bernardo do Campo (SG). In addition, we considered specimens deposited in the Museu Nacional, Rio de Janeiro (MNRJ), mentioned in the literature. We did not use published records



**FIGURE 1:** Map showing mountaintop regions of *campos rupestres* and *campos de altitude* considered in this review. Localities codes are presented in Table 1.

**TABLE 1:** Mountaintop regions of *campos rupestres* and *campos de altitude* considered in this review. Localities are listed from north to south and are grouped by Roman-Arabic number codes following the mountain ranges of each one: I = Espinhaço Range; II = Serra da Mantiqueira; III = Serra do Mar. Numerical codes for sources are presented in Appendix 1.

Code	Locality	Coordinates	Elevation (m)		Sources
I.1	Dias Coelho	11°29'S, 40°49'W	950	21	
I.2	Izabel Dias	11°30'S, 41°00'W	850	21	
I.3	Morro do Chapéu	11°34'S, 41°07'W	930-1,000	21, 23, 24, 31, 36, 38, 41, 49, 50, 58, 63, 68, 70, 71, 104, 107, 109, 151	
I.4	Paramirim	12°01'S, 42°15'W	1,050-1,280	21, 24, 68	
I.5	Palmeiras-Lençóis	Between 12°26'S, 41°25'W and 12°29'S, 41°30'W	900-1,250	21, 22, 23, 24, 50, 68, 70, 71, 82, 85, 87, 101, 104, 106, 132, 167	
I.6	Serra do Ribeirão	12°33'S, 41°25'W	950	132	
I.7	Vale do Paty	12°48'S, 41°25'W	1,035	132	
I.8	Serra de Andaraí	Between 12°46'S, 41°19'W and 12°51'S, 41°23'W	730-1,150	21, 22, 23, 24, 33, 50, 58, 68, 151, 157	
I.9	Igatu	Between 12°52'S, 41°18'W and 12°53'S, 41°20'W	730-950	21, 22, 23, 24, 29, 30, 50, 68, 132, 159	
I.10	Mucugê	Between 12°57'S, 41°20'W and 13°01'S, 41°23'W	870-1,320	21, 22, 23, 24, 33, 50, 51, 68, 70, 71, 82, 92, 96, 97, 98, 99, 100, 104, 106, 113, 126, 132, 134, 135, 152, 157, 158, 159, 167	
I.11	Roncador	Between 13°08'S, 41°21'W and 13°08'S, 41°22'W	1,000-1,490	21	
I.12	Piatã	Between 13°08'S, 41°44'W and 13°10'S, 41°46'W	1,220-1,640	21, 24, 50, 68	
I.13	Serra do Cocal	Between 13°14'S, 41°44'W and 13°16'S, 41°45'W	900-1,170	21	
I.14	Morro do Ouro	Between 13°15'S, 42°11'W and 13°16'S, 42°11'W	900-1,120	21	
I.15	Serra das Almas	Between 13°31'S, 41°49'W and 13°35'S, 41°58'W	980-1,950	68, 70, 71, 82, 104, 157, 167	
I.16	Espigão do Taquari	13°36'S, 41°27'W	1,130-1,235	21	
I.17	Barra da Estiva	Between 13°40'S, 41°16'W and 13°41'S, 41°18'W	1,000-1,500	21, 23, 24, 50, 58, 68, 107	
I.18	Brejinho das Ametistas	14°21'S, 42°32'W	1,050	152, 167	
I.19	Jacaraci	14°53'S, 42°31'W	1,330	167	
I.20	Serra do Pau D'Arco	Between 15°03'S, 42°40'W and 15°04'S, 42°41'W	1,150-1,600	104, 140, 152, 167	
I.21	Serra da Formosa	Between 15°12'S, 42°47'W and 15°15'S, 42°50'W	1,000-1,780	144, 152, 167	
I.22	Campos Geraes	15°27'S, 42°45'W	1,200	2, 151	
I.23	Gerais de Santana	Between 15°41'S, 42°48'W and 15°44'S, 42°50'W	1,100-1,400	144, 167	
I.24	Serra do Barão	Between 16°12'S, 42°52'W and 16°36'S, 42°56'W	800-1,250	21, 23, 24, 25, 27, 38, 50, 58, 63, 104, 107, 139, 153, 167	
I.25	Campina do Bananal	Between 16°50'S, 43°01'W and 16°51'S, 43°03'W	1,100-1,360	94, 104, 107, 139, 140, 150, 152, 156, 167	
I.26	Serra Resplandecente	Between 16°59'S, 43°19'W and 17°02'S, 43°21'W	1,020-1,320	36, 104, 139, 150, 152, 167	
I.27	Serra do Cabral	Between 17°41'S, 44°10'W and 17°45'S, 44°17'W	950-1,250	127, 140, 152, 167	
I.28	Serra dos Poções	18°04'S, 43°25'W	980	36, 38, 63, 104	
I.29	Serra do Ambrósio	Between 18°06'S, 43°02'W and 18°08'S, 43°08'W	880-1,520	152	
I.30	Diamantina	Between 18°08'S, 43°31'W and 18°16'S, 43°38'W	1,000-1,370	1, 6, 17, 18, 21, 23, 24, 25, 27, 36, 37, 38, 50, 58, 104, 107, 151, 152, 153, 157, 159, 167	
I.31	Datas	Between 18°23'S, 43°36'W and 18°27'S, 43°39'W	1,020-1,370	36, 104	
I.32	Serra do Gavião	Between 18°09'S, 43°17'W and 18°32'S, 43°27'W	1,000-2,000	38, 62, 63, 74, 93, 94, 104, 107, 108, 127, 129, 150, 152, 167	

TABLE 1: Continued.

Code	Locality	Coordinates	Elevation (m)	Sources
I.33	Serra do Barro Preto	Between 18°35'S, 43°53'W and 18°39'S, 43°55'W	1,170-1,300	150, 152, 167
I.34	Serra Talhada	18°48'S, 43°53'W	1,000-1,035	62, 94
I.35	Serra do Cipó	Between 19°02'S, 43°25'W and 19°25'S, 43°43'W	1,150-1,690	21, 23, 24, 29, 30, 34, 36, 37, 38, 41, 42, 44, 46, 47, 48, 49, 50, 57, 58, 61, 62, 63, 74, 84, 87, 89, 91, 94, 104, 107, 110, 111, 112, 114, 115, 116, 117, 118, 119, 125, 131, 140, 142, 144, 145, 146, 147, 149, 150, 151, 152, 153, 157, 159, 160, 167
I.36	Serra da Piedade	Between 19°48'S, 43°40'W and 19°49'S, 43°41'W	1,400-1,740	18, 21, 24, 37, 50, 58, 63, 74, 77, 89, 95, 102, 104, 107, 140, 152, 153, 167
I.37	Serra da Água Limpa	Between 19°49'S, 43°30'W and 19°54'S, 43°31'W	1,000-1,400	161, 167
I.38	Serra do Curral	Between 19°57'S, 43°54'W and 19°58'S, 43°55'W	1,250-1,330	53, 59, 63, 66, 67, 74, 75, 76, 77, 87, 89, 104, 138, 167
I.39	Serra do Rola-Moça	Between 20°00'S, 43°58'W and 20°05'S, 44°00'W	1,300-1,450	77, 87, 104, 105, 120, 121, 122, 123, 128, 133, 152, 154, 167
I.40	Serra da Gandarela	Between 20°03'S, 43°39'W and 20°06'S, 43°42'W	1,470-1,640	38, 58, 63, 104, 152, 164, 167
I.41	Serra de Itatiaiuçu	Between 20°07'S, 44°19'W and 20°07'S, 44°21'W	1,230-1,340	51, 159
I.42	Serra do Caraça	Between 20°03'S, 43°26'W and 20°08'S, 43°31'W	1,200-2,070	6, 17, 21, 24, 29, 30, 35, 36, 37, 38, 40, 50, 51, 58, 63, 64, 74, 81, 86, 87, 88, 89, 90, 102, 103, 104, 107, 108, 129, 141, 149, 151, 152, 153, 155, 157, 158, 159, 160, 167
I.43	Alegria	Between 20°07'S, 43°25'W and 20°12'S, 43°23'W	900-1,000	24, 50, 107, 151, 152, 153, 167
I.44	Serra do Capanema	Between 20°10'S, 43°36'W and 20°11'S, 43°37'W	1,400-1,700	104, 152, 163, 167
I.45	Serra Santa	Between 20°11'S, 43°50'W and 20°14'S, 43°52'W	1,110-1,350	18, 24, 29, 30, 104, 107, 158, 159, 160
I.46	Serra do Batatal	Between 20°13'S, 43°30'W and 20°18'S, 43°34'W	1,240-1,810	38, 102, 104, 152, 157, 167
I.47	Serra da Moeda	Between 20°13'S, 43°56'W and 20°22'S, 43°58'W	1,270-1,480	38, 58, 63, 74, 77, 89, 104, 107, 167
I.48	Pico do Itacolomi	Between 20°24'S, 43°30'W and 20°26'S, 43°26'W	1,200-1,720	6, 17, 18, 21, 23, 24, 37, 38, 60, 77, 87, 89, 102, 104, 167
I.49	Serra de Belo Vale	Between 20°26'S, 43°56'W and 20°27'S, 43°56'W	1,260-1,520	77, 152
I.50	Serra do Mascate	Between 20°27'S, 43°55'W and 20°28'S, 43°56'W	1,350-1,620	152, 167
I.51	Serra de Ouro Branco	Between 20°28'S, 43°35'W and 20°30'S, 43°44'W	1,150-1,540	38, 63, 104, 107, 157, 167
II.1	Serra do Caparaó	Between 20°22'S, 41°47'W and 20°29'S, 41°50'W	1,900-2,890	10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 26, 28, 32, 45, 50, 58, 64, 65, 69, 78, 83, 102, 104, 130, 151, 167
II.2	Serra da Vargem Grande	20°30'S, 42°04'W	1,500-1,750	140, 152, 167
II.3	Serra do Brigadeiro	Between 20°33'S, 42°25'W and 20°47'S, 42°28'W	1,300-1,980	72
II.4	Serra de São José	Between 21°02'S, 44°06'W and 21°06'S, 44°13'W	1,000-1,320	63, 87, 104
II.5	Serra do Lenheiro	Between 21°08'S, 44°17'W and 21°09'S, 44°19'W	1,050-1,200	63, 104, 140, 167
II.6	Serra do Ibitipoca	Between 21°40'S, 43°52'W and 21°42'S, 43°54'W	1,400-1,780	54, 55, 104, 148, 152, 167
II.7	Serra do Papagaio	Between 22°00'S, 44°38'W and 22°03'S, 44°41'W	1,500-2,150	73, 104, 140, 143, 167
II.8	Serra do Itatiaia	Between 22°21'S, 44°38'W and 22°25'S, 44°43'W	2,000-2,787	3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 16, 17, 18, 19, 20, 26, 32, 38, 39, 43, 51, 52, 58, 64, 102, 104, 151, 157, 159, 167

TABLE 1: Continued.

Code	Locality	Coordinates	Elevation (m)		Sources
II.9	Pedra da Mina	22°25'S, 44°50'W	2,797	79	
II.10	Marins-Itaguapé	Between 22°29'S, 45°04'W and 22°29'S, 45°07'W	2,050-2,250	137, 156	
II.11	Pedra de São Domingos	22°41'S, 45°57'W	2,000	140, 152, 167	
II.12	Serra dos Poncianos	Between 22°53'S, 46°00'W and 22°53'S, 46°03'W	1,800-2,010	167	
II.13	Atibaia	Between 23°08'S, 46°30'W and 23°10'S, 46°31'W	1,050-1,350	56	
III.1	Pedra Azul	Between 20°24'S, 41°00'W and 20°24'S, 41°01'W	1,450-1,900	65, 80, 167	
III.2	Serra dos Órgãos	Between 22°20'S, 42°34'W and 22°29'S, 43°06'W	1,400-2,263	8, 11, 20, 26, 32, 52, 58, 64, 124, 136, 151, 157, 162, 165, 166, 167	
III.3	Serra da Bocaina	Between 22°42'S, 44°34'W and 22°49'S, 44°46'W	1,500-2,200	49, 52, 58, 104, 158, 159	

by Ihering (1900) and Ruschi (1978, 1982b), since those authors did not mention the elevation and habitats where they recorded the species (*see* Remsen, 1994). Based on our field experience, we also did not include species only recorded in some published checklists not supported by physical evidence (specimens, photos or tape-recordings) since these records are possibly based on misidentifications. A list of all sources used in this revision is presented in Appendix 1.

Geographical distribution patterns for bird species were discussed based on: 1) our own fieldwork conducted in the following regions: Brazil (Pará, Maranhão, Ceará, Pernambuco, Tocantins, Bahia, Goiás, Mato Grosso, Mato Grosso do Sul, Minas Gerais, Espírito Santo, Rio de Janeiro, São Paulo, Paraná, Santa Catarina, Rio Grande do Sul), Peru (Ancash), Bolivia (Santa Cruz), and Argentina (Misiones); 2) specimens housed in the above mentioned institutions; and 3) literature review (*see* Appendix 1).

Taxonomy follows the Brazilian Committee of Ornithological Records (CBRO, 2007). Bird species were classified following their endemism after several authors (Ridgely & Tudor, 1989, 1994; Silva, 1995, 1997; Stotz *et al.*, 1996; Sick, 1997; Stattersfield *et al.*, 1998; Brooks *et al.*, 1999; Silva & Bates, 2002; Silva & Santos, 2005). The exceptions are those typical species of the *campos rupestres* considered restricted to the Cerrado region (Silva, 1995; 1997; Silva & Bates, 2002; Silva & Santos, 2005). Here, they were considered endemics to the open-habitats of southeastern Brazilian mountaintops (following arguments presented by Vasconcelos *et al.*, 2003; Lopes, 2008; Vasconcelos, 2008b). These species are *Augastes scutatus*, *Asthenes luizae*, *Polystictus superciliaris*, and *Embernagra longicauda*.

## RESULTS AND DISCUSSION

### Species richness and composition

A total of 231 bird species was recorded in the open-habitats of southeastern Brazilian mountaintops (Appendix 2). The list for the *campos rupestres* has a total of 205 species, while the *campos de altitude* has 123 species (Appendix 2). Ninety-seven species occur both in the *campos rupestres* and in the *campos de altitude*. One hundred and eight species were exclusive to the *campos rupestres*, while 26 species were recorded only in the *campos de altitude* (Appendix 2).

The higher species richness recorded in the *campos rupestres* can be attributed to the Espinhaço Range being in the contact zones among Atlantic Forest, Cerrado, and Caatinga (Giulietti & Pirani, 1988; Ab'Sáber, 1990; Harley, 1995; Giulietti *et al.*, 1997). Such a variety of biomes within the Espinhaço Range leads to an increase in its species richness, especially because birds typical of the open-habitats of the Cerrado and semi-open-habitats of the Caatinga can also live in the *campos rupestres* (*see* Parrini *et al.*, 1999; Vasconcelos & D'Angelo-Neto, 2007). By contrast, the *campos de altitude* can be considered as isolated islands of open-habitat among forested areas in the Atlantic Forest region. Since many forest bird species cannot live in open areas, colonization of the *campos de altitude* by species from nearby habitats is more difficult than in the *campos rupestres*. Low temperature, episodic frosts and even snow (Sick, 1970; Martinelli & Orleans e Bragança, 1996; Safford, 1999b) may also preclude permanent colonization by open-habitat birds found in lower altitudes that could use current man-made pastures to get there. Furthermore, *campos rupestres* areas are more naturally interconnected compared with the *campos de altitude*.

This island feature of the *campos de altitude* could explain their lower species richness (MacArthur & Wilson, 1967). Meanwhile, more systematic surveys and area estimation based on satellite images are still necessary in order to test such hypothesis, as has been done for the northern Andes (Vuilleumier, 1970), Tepuis (Cook, 1974), and Sierras Pampeanas (Nores, 1995).

The differences in bird species composition between the *campos rupestres* and the *campos de altitude*, despite some species being common to both, demonstrate a pattern similar to that recognized for the flora (Safford, 1999a, 2007).

### Geographic distribution patterns of the avifauna

Five patterns of distribution among the bird species occurring in the *campos rupestres* and *campos de altitude* can be recognized (Table 2):

- Non-endemic (191 species): Widespread species non-endemic to any biome.
- Atlantic Forest endemics (26 species): Species endemic to the Atlantic Forest region in southeastern Brazil and adjacent areas (eastern Paraguay, northeastern Argentina, and Uruguay).
- Cerrado endemics (6 species): Species restricted to the Cerrado region, except those typical of the *campos rupestres*, included here in another category (see materials and methods).
- Caatinga endemic (1 species): Species endemic to the Caatinga region in northeastern Brazil.
- Montane open-habitat endemics (7 species): Species endemic to the open-habitats of southeastern Brazilian mountaintops (*campos rupestres* and *campos de altitude*), including those restricted to *campos rupestres* (see above).

The highest proportion of the species living in southeastern Brazilian montane open-habitats is

represented by widespread and non-endemic species (82.7%), with only 3.0% endemic to these habitats (Table 2). This is the opposite pattern of that found in the Andean puna and páramo, where 29% of the species are endemics and only 13.5% are widespread species (Vuilleumier, 1986). The high percentage of widespread bird species living in Brazilian montane open-habitats also contrasts to the local flora, which contains many endemic species (Eiten, 1992; Martinelli & Orleans e Bragança, 1996; Giulietti *et al.*, 1997; Safford, 1999a; Menezes & Giulietti, 2000).

The distribution patterns of the birds from *campos rupestres* and *campos de altitude* show remarkable differences between both vegetation types. Although both formations show a dominance of wide-ranging species, their percentage is a little higher in the *campos rupestres* (85.9%) than in the *campos de altitude* (79.7%) (Table 2).

Atlantic Forest birds from southeastern Brazil and adjacent areas account for more species in the *campos de altitude* (17.1%) than in the *campos rupestres* (7.3%) (Table 2). The *campos de altitude* have 21 of the 26 Atlantic Forest endemics (80.8%) that have been recorded in the montane open-habitats of southeastern Brazil, while in the *campos rupestres*, there are only 15 (57.7%) (Table 2). The ranges of many Atlantic Forest species found in the *campos de altitude* do not reach northwest to the *campos rupestres* region. Some examples are: *Stephanoxis lalandi*, *Drymophila genei*, *Scytalopus notorius*, *Hemitriccus obsoletus*, *Phylloscartes difficilis*, *Poospiza thoracica*, and *Saltator maxillosus*. Nevertheless, some Atlantic Forest endemics do reach the *campos rupestres*, especially at localities in the southern Espinhaço Range (Vasconcelos *et al.*, 1999b; Vasconcelos & Melo-Júnior, 2001) and their presence can be explained by the geographical proximity of this region to the coastal ranges. Such taxa include: *Leucochloris albicollis*, *Muscipira vetula*, *Tangara desmaresti*, and *Haplospiza unicolor*.

**TABLE 2:** Distribution of bird species of *campos rupestres* and *campos de altitude*, analyzed as a single unit and separately. Distribution pattern: NE = non-endemic; AF = endemic to the Atlantic Forest region; CA = endemic to the Caatinga region; CE = endemic to the Cerrado region; MT = endemic to the open-habitats of southeastern Brazilian mountaintops (*campos rupestres* and *campos de altitude*).

Distribution pattern	<i>Campos rupestres</i> and <i>campos de altitude</i>		<i>Campos rupestres</i>		<i>Campos de altitude</i>	
	N	%	N	%	N	%
NE	191	82.7	176	85.9	98	79.7
AF	26	11.3	15	7.3	21	17.1
CE	6	2.6	6	2.9	1	0.8
CA	1	0.4	1	0.5	0	0
MT	7	3.0	7	3.4	3	2.4
Total	231	100	205	100	123	100

Among the bird species restricted to the Atlantic Forest region, four show a distribution pattern associated with the highlands in southeastern Brazil, despite the fact that they also occur either in lowlands or highlands in subtropical regions of southern Brazil and adjacent areas (south of 23°30'S). Those species are: *Stephanoxis lalandi*, *Scytalopus iraiensis*, *Stephanophorus diadematus*, and *Emberizoides ypiranganus*.

This pattern was also recognized in some plant genera (Simpson, 1979; Safford, 1999a), bees (Silveira & Cure, 1993), and a rodent (Gonçalves *et al.*, 2007), suggesting that these taxa probably had larger ranges during the Pleistocene (Simpson, 1979; Silveira & Cure, 1993; Safford, 1999a, 2007; Gonçalves *et al.*, 2007). During interglacial periods, with the predominance of warmer climates, the ranges of these taxa would have retreated southward and upward, with the probable trapping of several taxa in the "glacial refugia" provided by the southeastern Brazilian mountaintops. In the case of *S. diadematus*, populations do not appear to be isolated, with possible connections throughout coastal São Paulo. Nevertheless, *S. lalandi*, *S. iraiensis*, and *E. ypiranganus* appear to have two main isolated population centers, one in the southeastern Brazilian mountains and another in forests and wetlands south of 23°30'S. We could not find any geographic variation in specimens of *S. iraiensis* and *E. ypiranganus* we collected in southeastern Brazilian highlands compared to southern specimens (see Vasconcelos *et al.*, 2006, 2008a). Forest expansion over grasslands in southeastern Brazil appears to be recent (Behling, 2002) and this could explain the absence of morphological differentiation in those species.

Nevertheless, *S. lalandi* is represented by two very different subspecies: *S. l. lalandi* (southeastern Brazilian mountains) and *S. l. loddigesii* (lowlands and/or highlands south of 23°30'S). Since these geographical forms are morphologically distinctive, do not have any known intermediate specimen, and do not present any known range overlap, it is suggested that each one can be considered a valid species.

This same pattern of distribution is shown by *Poospiza lateralis* / *P. cabanisi* (Assis *et al.*, 2007). The pattern found for *S. lalandi* and *P. lateralis* groups suggests that the original ranges of the ancestral taxa have retreated to two areas with cooler and moister climatic features, with subsequent isolation and differentiation: 1) the high mountains of coastal southeastern Brazil; and 2) southern Brazil and adjacent areas (eastern Paraguay, northeastern Argentina, and Uruguay). This pattern should also be associated to Plio-Pleistocene tectonic movements (Petri & Fúlvaro, 1986) that

also appear to have influenced the differentiation of some forest bird species (Pinto, 1978; Silva & Stotz, 1992; Silva & Straube, 1996).

Another southern montane species recently found in the southeastern Brazilian mountaintops is *Cinclodes pabsti* (Freitas *et al.*, 2008). Nevertheless, it is necessary to analyze more specimens and voice recordings of this northern population to assess if it can be treated as a new taxon.

No species endemic to the Caatinga was recorded in the *campos de altitude* and only one Cerrado endemic (0.8%) was found in this habitat type. Small percentages of Cerrado (2.9%) and Caatinga endemics (0.5%) were found among the *campos rupestres* avifauna (Table 2).

All Cerrado endemics recorded in the *campos rupestres* have original distributions that do not reach the coastal ranges, except *Cyanocorax cristatellus*. This species have been expanding its range eastward and southward due to the deforestation of the Atlantic Forest (Alvarenga, 1990; Vasconcelos, 1999a; Mallet-Rodrigues *et al.*, 2007; Lopes, 2008), but it is possible that it is a native species in the only area of *campos de altitude* it was recorded (Serra do Ibitipoca), since the region is in a transitional zone between Atlantic Forest and Cerrado, with several bird species typical of each biome (Pacheco *et al.*, 2008).

The Caatinga reaches the borders of the central and northern Espinhaço Range, what explains the occurrence of one of its endemics (*Sakesphorus cristatus*) in the *campos rupestres*. Despite the fact that Marini & Lopes (2005) found *S. cristatus* in the Rio Doce basin at Conceição do Mato Dentro, we still consider it as a Caatinga endemic since it is probably expanding its range due to deforestation, as observed for other typical Caatinga birds in this river valley, such as *Columbina picui*, *Compsothraupis loricata*, and *Icterus jamacaii* (M.F.V., *pers. obs.*).

Bird species that have ranges restricted to mountaintop open-habitats of southeastern Brazil are represented by low percentages in both formations: 3.4% in the *campos rupestres* and 2.4% in the *campos de altitude* (Table 2). Montane open-habitat endemics with ranges restricted to the *campos rupestres* of the Espinhaço Range are: *Augastes lumachella*, *Augastes scutatus*, *Formicivora grantsau*, and *Asthenes luizae*. *A. lumachella* and *F. grantsau* are endemic to the northern Espinhaço Range, while *A. scutatus* and *A. luizae* are restricted to the central and southern regions (Vasconcelos, 2008b). *Oreophylax moreirae* was considered restricted to the *campos de altitude* in coastal mountains (Serra da Mantiqueira, Serra dos Órgãos, and Serra do Caparaó), but was recently discovered in the highest



peaks of the southern Espinhaço Range, at Serra do Caraça (Melo-Júnior *et al.*, 1998; Vasconcelos, 2000; Vasconcelos & Melo-Júnior, 2001). *Polystictus superciliaris* and *Embernagra longicauda* were considered endemic to the *campos rupestres* of the Espinhaço Range, and also as Cerrado endemics (Silva, 1995, 1997; Silva & Bates, 2002; Silva & Santos, 2005). Nevertheless, these species were recorded outside the Espinhaço Range, in the Serra da Canastra and in the coastal mountains of the Atlantic Forest region, such as Serra do Mar, Serra da Mantiqueira, and Serra do Caparaó (Sick, 1997; Silveira, 1998; Vasconcelos, 1999a, b, 2003, 2008a, b; Vasconcelos *et al.*, 2003). The occurrence of *E. longicauda*, a typical species of the serras in the Espinhaço Range, in the Caparaó highlands suggests past connections between the *campos de altitude* of this area and the intermost open areas of Central Brazil, a pattern also corroborated by a recent study on *Akodon* rodents (Gonçalves *et al.*, 2007). Therefore, among the seven endemic bird species of the open-habitats of southeastern Brazilian mountaintops, four are restricted to the *campos rupestres* of the Espinhaço Range, but none of them is restricted to the *campos de altitude* of the eastern mountains.

In addition to those endemic species, there are four subspecies that appear to be restricted to the *campos rupestres* area of the Espinhaço Range: *Phaethornis pretrei schwarti*, *Campylopterus largipennis diamantiniensis*, *Colibri delphinae greenewalti*, and *Knipolegus nigerrimus hoflingi*. Nevertheless, validity of some of those taxa are questionable (Vielliard, 1994; Stiles, 1999; Brammer, 2002) and further taxonomic studies are still necessary.

The majority of birds living in the open-habitats of southeastern Brazilian mountaintops are represented by widespread and non-endemic species. Thus, the Andean-Patagonian influence in the avifauna in those habitats is small. Nevertheless, four endemic species appear to have close relationships with other Andean (*Augastes lumachella*, *A. scutatus*, and *Oreophylax moreirae*) and Andean-Chacoan-Patagonian (*Asthenes luizae*) species (Sick, 1970, 1985; Silva, 1995; Vielliard, 1996; Vasconcelos *et al.*, 2008b), but in the absence of any phylogenetic analysis, it is impossible to determine the precise evolutionary relationships and biogeographical origins of these species.

### Conservation

This review shows that bird species endemic to the Atlantic Forest, Cerrado, and Caatinga may

be found together at the montane open-habitats of southeastern Brazil. Besides this contact zone character, those habitats have their own few endemic birds. Among the birds of the montane open-habitats of southeastern Brazil, 16 are considered threatened or near-threatened either globally or in Brazil (Table 3). There are some discrepancies between both checklists. For instance, in Brazil, there are four threatened and two near-threatened species (Machado *et al.*, 2005), while globally there are six threatened and 10 near-threatened species (BirdLife International, 2007) (Table 3). Two examples are *Asthenes luizae* and *Poospiza cinerea*, both not considered as threatened species in Brazil, but classified as globally vulnerable (Table 3).

Despite 32 protected areas established in *campos rupestres* and *campos de altitude*, especially national and state parks (Table 4), those reserves cannot guarantee the conservation of all of their vegetation types and endemic and threatened birds found atop those mountain ranges. The first reason is, following a rule in Brazil and other developing countries, many of the reserves exist only “on paper”, and there is neither financial support nor control of illegal activities

**TABLE 3:** Bird species considered threatened or near-threatened in Brazil and in the world. Status: EN = endangered; NT = near-threatened; VU = vulnerable.

Family/Species	Brazil	Global
Tinamidae		
<i>Nothura minor</i>	VU	VU
Accipitridae		
<i>Harpyhaliaetus coronatus</i>	VU	EN
Psittacidae		
<i>Primolius maracana</i>	NT	NT
<i>Aratinga auricapillus</i>	—	NT
Trochilidae		
<i>Augastes scutatus</i>	—	NT
<i>Augastes lumachella</i>	—	NT
Thamnophilidae		
<i>Drymophila genei</i>	—	NT
Rhinocryptidae		
<i>Scytalopus iraiensis</i>	EN	EN
Furnariidae		
<i>Cinclodes pabsti</i>	NT	NT
<i>Asthenes luizae</i>	—	VU
Tyrannidae		
<i>Polystictus superciliaris</i>	—	NT
<i>Phylloscartes difficilis</i>	—	NT
<i>Culicivora caudacuta</i>	VU	VU
Emberizidae		
<i>Porphyospiza caerulescens</i>	—	NT
<i>Poospiza cinerea</i>	—	VU
<i>Embernagra longicauda</i>	—	NT

**TABLE 4:** Reserves in the localities of *campos rupestres* and *campos de altitude* analyzed in this study. Numerical codes for localities are presented in Table 1. Brazilian states: BA = Bahia; ES = Espírito Santo; MG = Minas Gerais; RJ = Rio de Janeiro; SP = São Paulo.

Reserves	States	Localities
Morro do Chapéu State Park, Monumento Natural Cachoeira do Ferro Doido	BA	I.3
Chapada Diamantina National Park, Mucugê Municipal Park	BA	I.5, I.6, I.7, I.8, I.9, I.10
Serra das Almas Municipal Park, Área de Relevante Interesse Ecológico Nascentes do Rio de Contas	BA	I.12, I.15
Serra Nova State Park	MG	I.23
Grão Mogol State Park	MG	I.24
Serra do Cabral State Park	MG	I.27
Biribiri State Park	MG	I.30
Pico do Itambé State Park, Rio Preto State Park	MG	I.32
Serra do Cipó National Park, Serra do Intendente State Park, Salão de Pedras Natural Park, Monumento Natural Serra da Ferrugem	MG	I.35
Monumento Natural Serra da Piedade	MG	I.36
Mangabeiras Municipal Park	MG	I.38
Serra do Rola Moça State Park	MG	I.39
Reserva Particular do Patrimônio Natural do Caraça	MG	I.42
Itacolomi State Park	MG	I.48
Serra do Caparaó National Park	ES/MG	II.1
Serra do Brigadeiro State Park	MG	II.3
Ibitipoca State Park	MG	II.6
Serra do Papagaio State Park	MG	II.7
Itatiaia National Park	MG/RJ	II.8
Itapetinga Grota Funda Municipal Park	SP	II.13
Pedra Azul State Park	ES	III.1
Serra dos Órgãos National Park, Desengano State Park, Três Picos State Park	RJ	III.2
Serra da Bocaina National Park	RJ/SP	III.3

inside their areas. Examples of paper parks with little enforcement are Serra Nova State Park, Grão Mogol State Park, Serra do Cabral State Park, and Sempre Vivas National Park (all in Minas Gerais state), the last one not included in this revision, but in the main massif of Espinhaço Range with large areas of *campos rupestres*. Most of these reserves were decreed without local landowners receiving compensation for their lands. Besides lack of funds, the main reasons for that are the lack of legal titles by the *de facto* owners and the generalized mess of land tenure in Brazil. As a result, there were reports of landowners setting fire to the *campos rupestres* of Serra do Cabral and Grão Mogol State Parks as a way of demanding the money they never received (*see below*).

Secondly, the staff working in most Brazilian protected areas consists of too few people unable to control and protect large areas against the pillaging of the natural heritage. Also, most reserves are threatened by uncontrolled tourism in southeastern Brazil. The recently created “Estrada Real” touristic route linking the southeastern Brazilian coast (Parati and Rio de Janeiro) to the highlands of Minas Gerais state, is an important threat to the conservation of the *campos rupestres* and *campos de altitude* as the number

of tourists visiting these mountainous areas has shown a rapid increase in the last three years without proper control on their activities. A portion of this “ecological road” partially destroyed the *campos rupestres* in the type-locality of *Asthenes luizae*, a globally vulnerable species (BirdLife International, 2007). This kind of development is a serious threat to this endemic bird species, as well as to a restricted-range plant species (Viana *et al.*, 2005). In the *campos de altitude* of Caparaó, Itatiaia, and Serra dos Órgãos National Parks, as well as in the Ibitipoca State Park, large numbers of tourists camp in the mountaintops, trampling the fragile native vegetation and causing fire hazards. It is common to find hundreds of people in those areas during weekends, holidays, or vacations. In Caparaó National Park, mules take the tourists to the highlands of this important area (Vasconcelos, 2003). These animals crush and graze the flora of the *campos de altitude* of Serra do Caparaó, composed of endemic and rare plant species (Leoni, 1997). So, management of these reserves is poor and obviously not conservation-focused. Additionally, some tourists also collect endemic plants like orchids and bromeliads (Conceição, 2000; Vasconcelos, 2000). In Reserva Particular do Patrimônio Natural do Caraça, a private

reserve, tourists used to collect endemic “canelas-de-ema” (Velloziaceae) in the *campos rupestres* as firewood (Vasconcelos, 2000).

Fire is another significant problem closely related to uncontrolled touristic activities, and large areas of Chapada Diamantina burn every year (Conceição, 2000). In the Serra do Caraça region, criminal fires affected one of the marshes where *Scytalopus iraiensis*, a globally endangered species at global and national levels, was recorded. After the fire, the species was not recorded again in this marsh (M.F.V., pers. obs.).

Unfortunately, many fires in the montane open-habitats are accidentally caused by tourists who camp in those areas, by ranchers renewing their pastures, or by disgruntled landowners who have not been paid for the land expropriated by parks and other reserves. Environmental modifications caused by fire in the *campos rupestres* and *campos de altitude* include floristic impoverishment and later occurrence of invasive plant species, which compete for space and light with native and endemic ones (Martinelli & Orleans e Bragança, 1996; Safford, 1999a; Vasconcelos, 2000; Vasconcelos *et al.*, 2002b).

Other human impacts are widespread in the mountains, such as mining, hunting, and overgrazing by cattle and horses (Safford, 1999a; Vasconcelos, 2000, 2003; Vasconcelos *et al.*, 2002b). In the Quadrilátero Ferrífero, extensive areas of *campos rupestres* growing on rock outcrops rich in iron ore (“canga”) have been quickly destroyed for mining (Jacobi *et al.*, 2007; Versieux & Wendt, 2007). The demise of this special type of *campos rupestres* probably led *Augastes scutatus*, a near-threatened species, to local extinction in one locality of Quadrilátero Ferrífero (Vasconcelos, 1999b). In the Cipó Range and Chapada Diamantina region, many families live based on the collection and sale of “sempre-vivas”, endemic plants of the family Eriocaulaceae, which are valuable for ornamental purposes.

An important set of endemic and threatened bird species live in the *campos rupestres* and *campos de altitude*, and better efforts must be directed toward the better conservation of those important vegetation types. Safford (1999a) presented a series of actions that should be taken as a base for the conservation of the *campos de altitude* flora. Those can also be applied to avifauna conservation in the highlands of southeastern Brazil.

## RESUMO

*Os ambientes abertos das montanhas do sudeste do Brasil são representados pelos campos rupestres (principalmente*

*na Cadeia do Espinhaço) e pelos campos de altitude (nas Serras do Mar e da Mantiqueira). Apesar da ocorrência de espécies endêmicas em ambos os tipos de vegetação, uma análise e síntese de suas comunidades de aves nunca foram realizadas. Nesse estudo, apresentamos uma revisão da avifauna dessas áreas, descrevemos padrões de distribuição geográfica e comentamos sobre a conservação destes ambientes e de sua avifauna. Um total de 231 espécies de aves foi registrado nos ambientes abertos dos topos de montanha do sudeste do Brasil. Nos campos rupestres, 205 espécies foram registradas, enquanto nos campos de altitude o total foi de 123 espécies. Cinco padrões de distribuição são reconhecidos dentre as aves que ocorrem nestes habitats: não-endêmicas (191 espécies), endêmicas da Mata Atlântica (26 espécies), endêmicas do Cerrado (6 espécies), endêmica da Caatinga (1 espécie) e endêmicas dos habitats abertos dos topos de montanha (7 espécies). Apesar da presença de várias unidades de conservação nessas regiões, estas reservas não garantem a conservação de seus importantes tipos vegetacionais e de suas avifaunas sob os atuais níveis de gestão. Uma vez que várias espécies de aves endêmicas e ameaçadas de extinção vivem nos campos rupestres e campos de altitude, maiores esforços devem ser direcionados à sua conservação.*

PALAVRAS-CHAVE: Avifauna; Campos rupestres; Campos de altitude; Distribuição; Conservação.

## ACKNOWLEDGEMENTS

We thank Brehm Foundation, World Wildlife Fund (WWF), and United States Agency for International Development (USAID) for financial support during this study. M.F.V. thanks Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis (IBAMA) for the permission to collect bird specimens in Brazil, Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for a master and a doctoral fellowship, and AMNH for a Collection Study Grant to work on eastern Brazilian birds. M.R. is supported by a grant from CNPq (1998-2001, 2007-2009 and 473428/2004-9), Fundação O Boticário de Proteção à Natureza, and Fapemig (APQ-0434-5.03/07). D. Hoffmann kindly prepared the map. C.R.M. Abreu, C.E.A. Carvalho, E.P.M. Carvalho-Filho, S. D'Angelo-Neto, H.B. Gomes, E. Gouvêa (*in memoriam*), D. Hoffmann, J. Julio-Filho, M. Maldonado-Coelho, G.N. Maurício, T.A. Melo-Júnior, R. Nassau e Braga, and L.F. Silveira, helped in the field. M. Votaw, F.A. Silveira, K.-L. Schuchmann, and L.E. Lopes kindly commented on draft versions of the manuscript. F.F. Vasconcelos, F.

Mallet-Rodrigues, F.S. Leite, G.B. Malacco, J.F. Pacheco, and R. Parrini shared their records based on personal observations from some localities. We also thank the staffs of the museums visited during this study, especially: J. Cracraft, P. Sweet, and P. Hart (AMNH), M. Hoffmann (MBML), L. Guimarães (MCN), C.S. Fontana (MCP), H.M.F. Alvarenga (MHNT), A. Aleixo (MPEG), L.F. Silveira (MZUSP), and R. Grantsau (SG).

## REFERENCES

- AB'SÁBER, A.N. 1990. Paleoclimas Quaternários e pré-história da América tropical – II. *Revista Brasileira de Biologia*, 50:821-831.
- ABREU, C.R.M. 2006. Revisão taxonômica de *Augastes scutatus* (Temminck, 1824) (Aves: Trochilidae). (M.Sc. Dissertation). Universidade Federal do Rio de Janeiro, Rio de Janeiro.
- ALMEIDA, A.C.C. & RAPOSO, M.A. 1999. Aspectos da biologia e abundância do beija-flor-de-gravatinha-vermelha, *Augastes lumachellus* (Lesson, 1838) (Aves: Trochilidae), em Morro do Chapéu, BA. *Revista Nordestina de Biologia*, 13:69-85.
- ALVARENGA, H.M.F. 1990. Novos registros e expansões geográficas de aves no leste do estado de São Paulo. *Ararajuba*, 1:115-117.
- ALVES, A.C.F.; MOTA, N.F.O.; VIANA, P.L.; MARQUES, D.A.; MORAES, P.O. & SALINO, A. 2007. O banho de *Augastes scutatus* (Temminck, 1824) em duas localidades de campos rupestres em Minas Gerais. *Atualidades Ornitológicas*, 137:48-49.
- ALVES, A.C.F.; MOTA, N.F.O.; VIANA, P.L.; MORAES, P.O. & SALINO, A. 2006. Novas observações sobre o banho de *Augastes scutatus* (Aves: Trochilidae). In: Ribon, R. (Ed.), *Resumos do XIV Congresso Brasileiro de Ornitologia*. Universidade Federal de Ouro Preto, Ouro Preto, p.26.
- ALVES, R.J.V. & KOLBEK, J. 1994. Plant species endemism in savanna vegetation on table mountains (campo rupestre) in Brazil. *Vegetatio*, 113:125-139.
- ALVES, R.J.V. & KOLBEK, J. 2009. Summit vascular flora of Serra de São José, Minas Gerais, Brazil. *Check List*, 5:35-73.
- ALVES, R.J.V.; CARDIN, L. & KROPF, M.S. 2007. Angiosperm disjunction "campos rupestres – restingas": a re-evaluation. *Acta Botanica Brasílica*, 21:675-685.
- ANDRADE, M.A. 1997a. As aves na região do Parque Estadual do Ibitipoca: conservação e distribuição. In: Rocha, G.C. (Ed.), *Anais do 1º Seminário de Pesquisa sobre o Parque Estadual do Ibitipoca*. Núcleo de Pesquisa em Zoneamento Ambiental da Universidade Federal de Juiz de Fora, Juiz de Fora, p.61-72.
- ANDRADE, M.A. 1997b. O Parque do Ibitipoca e suas aves. *Uiraçu*, 1:5.
- ANDRADE, M.A. 1998. O Parque Estadual do Itacolomi e suas aves. *Uiraçu*, 2:4.
- ANDRADE, M.A.; ANDRADE, M.V.G.; GONTIJO, R.G.R. & SOUZA, P.O. 1998. Ocorrência do Cipó-canastero (*Asthenes luizae*) e do Gavião-pernilongo (*Geranospiza caerulescens*) no interior do Parque Nacional da Serra do Cipó, Minas Gerais. *Atualidades Ornitológicas*, 82:10.
- ANDRADE, P.M.; GONTIJO, T.A. & GRANDI, T.S.M. 1986. Composição florística e aspectos estruturais de uma área de "campo rupestre" do Morro do Chapéu, Nova Lima, Minas Gerais. *Revista Brasileira de Botânica*, 9:13-21.
- ASSIS, C.P.; RAPOSO, M.A. & PARRINI, R. 2007. Validação de *Poospiza cabanisi* Bonaparte, 1850 (Passeriformes: Emberizidae). *Revista Brasileira de Ornitologia*, 15:103-112.
- BAUER, C. 1999. *Padrões atuais de distribuição das aves florestais na região sul do estado do Espírito Santo, Brasil*. (M.Sc. Dissertation). Universidade Federal do Rio de Janeiro, Rio de Janeiro.
- BEHLING, H. 1998. Late Quaternary vegetational and climatic changes in Brazil. *Review of Palaeobotany and Palynology*, 99:143-156.
- BEHLING, H. 2002. South and southeast Brazilian grasslands during late Quaternary times: a synthesis. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 177:19-27.
- BENITES, V.M.; CAIAFA, A.N.; MENDONÇA, E.S.; SCHAEFER, C.E. & KER, J.C. 2003. Solos e vegetação nos complexos rupestres de altitude da Mantiqueira e do Espinhaço. *Floresta e Ambiente*, 10:76-85.
- BENITES, V.M.; SCHAEFER, C.E.G.R.; SIMAS, F.N.B. & SANTOS, H.G. 2007. Soils associated with rock outcrops in the Brazilian mountain ranges Mantiqueira and Espinhaço. *Revista Brasileira de Botânica*, 30:569-577.
- BIRDLIFE INTERNATIONAL. 2007. *Search for species*. Available at: <www.birdlife.org/datazone/species/index>. Access in: 31/10/2007.
- BRAMMER, F.P. 2002. *Species concepts and conservation priorities: a study of birds in north-east Brazil*. (M.Sc. Dissertation). University of Copenhagen, Copenhagen.
- BROOKS, T.; TOBIAS, J. & BALMFORD, A. 1999. Deforestation and bird extinctions in the Atlantic forest. *Animal Conservation*, 2:211-222.
- CAIAFA, A.N. & SILVA, A.F. 2005. Composição florística e espectro biológico de um campo de altitude no Parque Estadual da Serra do Brigadeiro, Minas Gerais, Brasil. *Rodriguésia*, 56:163-173.
- CAIAFA, A.N. & SILVA, A.F. 2007. Structural analysis of the vegetation on a highland granitic rock outcrop in southeast Brazil. *Revista Brasileira de Botânica*, 30:657-664.
- CARNEVALLI, N.E.D. 1980. Contribuição ao estudo da ornitofauna da Serra do Caraça, Minas Gerais. *Lundiana*, 1:89-98.
- CARNEVALLI, N.E.D. 1982. *Embernagra longicauda* Strikiland [sic], 1844; sua ocorrência em Minas Gerais, Brasil (Aves: Fringillidae). *Lundiana*, 2:85-88.
- CARVALHAES, A.M.P. 2001. *Dinâmica da comunidade de aves do Parque Nacional da Chapada Diamantina*. (Ph.D. Dissertation). Universidade Estadual Paulista, Botucatu.
- CBRO (COMITÊ BRASILEIRO DE REGISTROS ORNITOLÓGICOS). 2007. *Lista das aves do Brasil*. 6ª Edição. 16/08/2007. Available at: <www.cbro.org.br>. Access in: 21/08/2007.
- COELHO, A.G. & MACHADO, C.G. 2003. As espécies de beija-flores e seus recursos florais em uma área de campo rupestre da Chapada Diamantina, BA. In: Machado, C.G. (Ed.), *Resumos do XI Congresso Brasileiro de Ornitologia*. Universidade Estadual de Feira de Santana, Feira de Santana, p.11.
- COLAÇO, M.A.S.; FONSECA, R.B.S.; LAMBERT, S.M.; COSTA, C.B.N.; MACHADO, C.G. & BORBA, E.L. 2006. Biologia reprodutiva de *Melocactus glaucescens* Buining & Brederoo e *M. paucispinus* G. Heimen & R. Paul (Cactaceae), na Chapada Diamantina, nordeste do Brasil. *Revista Brasileira de Botânica*, 29:239-249.
- CONCEIÇÃO, A.A. & GIULIETTI, A.M. 2002. Composição florística e aspectos estruturais de campo rupestre em dois platôs do Morro do Pai Inácio, Chapada Diamantina, Bahia, Brasil. *Hoebnea*, 29:37-48.
- CONCEIÇÃO, A.A. & PIRANI, J.R. 2005. Delimitação de habitats em campos rupestres na Chapada Diamantina, Bahia: substratos, composição florística e aspectos estruturais. *Boletim de Botânica da Universidade de São Paulo*, 23:85-111.
- CONCEIÇÃO, A.A. 2000. Alerta para a conservação da biota na Chapada Diamantina. *Ciência Hoje*, 27:54-56.
- CONCEIÇÃO, A.A.; PIRANI, J.R. & MEIRELLES, S.T. 2007. Floristics, structure and soil of insular vegetation in four

- quartzite-sandstone outcrops of "Chapada Diamantina", northeast Brazil. *Revista Brasileira de Botânica*, 30:641-656.
- COOK, R.E. 1974. Origin of the highland avifauna of southern Venezuela. *Systematic Zoology*, 23:257-264.
- CORDEIRO, P.H.C.; MELO-JÚNIOR, T.A. & VASCONCELOS, M.F. 1998. A range extension for Cipó Canastero *Asthenes luizae* in Brazil. *Cotinga*, 10:64-65.
- COSTA, L.M. & RODRIGUES, M. 2006a. Estação reprodutiva de aves nos campos rupestres da Serra do Cipó, Minas Gerais. In: Ribon, R. (Ed.), *Resumos do XIV Congresso Brasileiro de Ornitologia*. Universidade Federal de Ouro Preto, Ouro Preto, p.24.
- COSTA, L.M. & RODRIGUES, M. 2006b. Área de vida e forrageamento de *Poospiza cinerea* (Aves: Emberezidae [sic]) na Serra do Cipó. In: Ribon, R. (Ed.), *Resumos do XIV Congresso Brasileiro de Ornitologia*. Universidade Federal de Ouro Preto, Ouro Preto, p.41.
- COSTA, L.M. & RODRIGUES, M. 2007. Novo registro de nidificação do beija-flor-de-gravata-verde *Augastes scutatus* estende período reprodutivo da espécie. *Revista Brasileira de Ornitologia*, 15:131-134.
- COSTA, L.M.; FREITAS, G.H.S.; RODRIGUES, M. & VASCONCELOS, M.F. 2008. New records of Lesser Grass Finch *Emberizoides ypiranganus* in Minas Gerais, Brazil. *Cotinga*, 29:182-183.
- DIAS, H.C.T.; FERNANDES-FILHO, E.I.; SCHAEFER, C.E.G.R.; FONTES, L.E.F. & VENTORIM, L.B. 2002. Geoambientes do Parque Estadual do Ibitipoca, município de Lima Duarte, MG. *Revista Árvore*, 26:777-786.
- DOMINGUES, L.A.L. & RODRIGUES, M. 2006. Territorialidade e área de vida de *Schistochlamys ruficapillus* (Aves: Thraupidae) na Serra do Cipó. In: Ribon, R. (Ed.), *Resumos do XIV Congresso Brasileiro de Ornitologia*. Universidade Federal de Ouro Preto, Ouro Preto, p.25.
- DORST, J. & VUILLEUMIER, F. 1986. Convergences in bird communities at high altitude in the tropics (especially the Andes and Africa) and at temperate latitudes (Tibet). In: Vuilleumier, F. & Monasterio, M. (Eds.), *High altitude tropical biogeography*. Oxford University Press, New York, p.120-149.
- EITEN, G. 1992. Natural Brazilian vegetation types and their causes. *Anais da Academia Brasileira de Ciências*, 64:35-65.
- FAUSTINO, T.C. & MACHADO, C.G. 2003. Os frutos consumidos por aves em uma área de campo rupestre da Chapada Diamantina, Bahia. In: Machado, C.G. (Ed.), *Resumos do XI Congresso Brasileiro de Ornitologia*. Universidade Estadual de Feira de Santana, Feira de Santana, p.14.
- FAUSTINO, T.C. & MACHADO, C.G. 2006. Frugivoria por aves em uma área de campo rupestre na Chapada Diamantina, BA. *Revista Brasileira de Ornitologia*, 14:137-143.
- FAUSTINO, T.C.; SANTANA, C.S. & MACHADO, C.G. 2003. As aves visitantes florais e frugívoras de *Hohenbergia ramageana* (Bromeliaceae) no Parque Municipal de Mucugê, Chapada Diamantina, BA. In: Machado, C.G. (Ed.), *Resumos do XI Congresso Brasileiro de Ornitologia*. Universidade Estadual de Feira de Santana, Feira de Santana, p.15.
- FREITAS, G.H.S. & RODRIGUES, M. 2008. Canário-rabudo *Embernagra longicauda*: uma ave típica das montanhas do leste do Brasil. *Atualidades Ornitológicas*, 141:20-21.
- FREITAS, G.H.S.; COSTA, L.M.; FERREIRA, J.D. & RODRIGUES, M. 2008. The range of Long-tailed Cinclodes *Cinclodes pabsti* extends to Minas Gerais (Brazil). *Bulletin of the British Ornithologists' Club*, 128:215-216.
- FUNCH, R.R. 1999. *A visitor's guide to the Chapada Diamantina mountains*. Secretaria de Ciência e Tecnologia, Salvador.
- GIULIETTI, A.M. & HENSOLD, N. 1990. Padrões de distribuição geográfica dos gêneros de Eriocaulaceae. *Acta Botanica Brasílica*, 4:133-158.
- GIULIETTI, A.M. & PIRANI, J.R. 1988. Patterns of geographic distribution of some plant species from the Espinhaço Range, Minas Gerais and Bahia, Brazil. In: Vanzolini, P.E. & Heyer, W.R. (Eds.), *Proceedings of a workshop on Neotropical distribution patterns*. Academia Brasileira de Ciências, Rio de Janeiro, p.39-69.
- GIULIETTI, A.M.; PIRANI, J.R. & HARLEY, R.M. 1997. Espinhaço Range region, eastern Brazil. In: Davis, S.D.; Heywood, V.H.; Herrera-MacBryde, O.; Villa-Lobos, J. & Hamilton, A.C. (Eds.), *Centres of plant diversity: a guide and strategy for their conservation*. Information Press, Oxford, v.3, p.397-404.
- GOMES, H.B. & GUERRA, T.J. 2006. Aves do Alto da Boa Vista, Serra do Cipó, Minas Gerais. In: Ribon, R. (Ed.), *Resumos do XIV Congresso Brasileiro de Ornitologia*. Universidade Federal de Ouro Preto, Ouro Preto, p.56.
- GOMES, H.B. & RODRIGUES, M. 2006a. Conservação de *Asthenes luizae* (Aves: Furnariidae) na Serra do Cipó, Minas Gerais. In: Ribon, R. (Ed.), *Resumos do XIV Congresso Brasileiro de Ornitologia*. Universidade Federal de Ouro Preto, Ouro Preto, p.32.
- GOMES, H.B. & RODRIGUES, M. 2006b. Padrão de canto de *Asthenes luizae* (Aves: Furnariidae) na Serra do Cipó, Minas Gerais. In: Ribon, R. (Ed.), *Resumos do XIV Congresso Brasileiro de Ornitologia*. Universidade Federal de Ouro Preto, Ouro Preto, p.43.
- GOMES, H.B. 2006. *Nidificação e padrão de canto de Asthenes luizae* (Aves: Furnariidae) na Serra do Cipó, Minas Gerais. (M.Sc. Dissertation). Universidade Federal de Minas Gerais, Belo Horizonte.
- GONÇALVES, P.R.; MYERS, P.; VILELA, J.F. & OLIVEIRA, J.A. 2007. Systematics of species of the genus *Akodon* (Rodentia: Sigmodontinae) in southeastern Brazil and implications for the biogeography of the campos de altitude. *Miscellaneous Publications Museum of Zoology, University of Michigan*, 197:1-24.
- GONZAGA, L.P. & CASTIGLIONI, G. 2001. *Aves das montanhas do sudeste do Brasil*. Sonopress, Manaus. CD-Audio.
- GONZAGA, L.P.; CARVALHAES, A.M.P. & BUZZETTI, D.R.C. 2007. A new species of *Formicivora* antwren from the Chapada Diamantina, eastern Brazil (Aves: Passeriformes: Thamnophilidae). *Zootaxa*, 1473:25-44.
- GOUNELLE, E. 1909. Contribution à l'étude de la distribution géographique des trochilidés dans le Brésil central et oriental. *Ornis*, 13:173-183.
- GRANTS AU, R. 1967. Sobre o gênero *Augastes*, com a descrição de uma subespécie nova (Aves, Trochilidae). *Papéis Avulsos de Zoologia*, 21:21-31.
- GRANTS AU, R. 1968. Die wiederentdeckung der brasilianischen kolibris *Augastes scutatus* und *Augastes lumachellus*. *Journal für Ornithologie*, 109:434-437.
- GRANTS AU, R. 1988. *Os beija-flores do Brasil*. Expressão e Cultura, Rio de Janeiro.
- GUERRA, T.J. & ALVES, A.C.F. 2006. Polinização e dispersão de sementes por aves em *Psittacanthus robustus* [sic] (Loranthaceae) em uma área de campo rupestre do sudeste brasileiro. In: Ribon, R. (Ed.), *Resumos do XIV Congresso Brasileiro de Ornitologia*. Universidade Federal de Ouro Preto, Ouro Preto, p.16.
- GUERRA, T.J.; GOMES, H.B. & GARCIA, F.I. 2006. Composição de bandos mistos em uma área de campo rupestre do sudeste brasileiro. In: Ribon, R. (Ed.), *Resumos do XIV Congresso Brasileiro de Ornitologia*. Universidade Federal de Ouro Preto, Ouro Preto, p.11.
- HARLEY, R.M. 1995. Introduction. In: Stannard, B.L.; Harvey, Y.B. & Harley, R.M. (Eds.), *Flora of the Pico das Almas*,

- Chapada Diamantina – Bahia, Brazil*. Royal Botanic Gardens, Kew, p.1-42.
- HEMMENDORFF, E. & MOREIRA, C. 1903. Relatório das excursões effectuadas na margem esquerda do Rio Branco em São Paulo e no Itatiaia na serra da Mantiqueira. *Archivos do Museu Nacional*, 12:159-167.
- HOFFMANN, D. & RODRIGUES, M. 2005. Táticas de forrageamento de *Polystictus superciliaris* (Aves: Tyrannidae), no Parque Estadual da Serra do Rola Moça, Nova Lima, MG. In: Aleixo, A. (Ed.), *Resumos do XIII Congresso Brasileiro de Ornitologia*. Universidade Federal do Pará, Belém, p.157.
- HOFFMANN, D. & RODRIGUES, M. 2006a. Biologia reprodutiva de *Polystictus superciliaris* (Aves: Tyrannidae) em área de canga no sudeste do Brasil. In: Ribon, R. (Ed.), *Resumos do XIV Congresso Brasileiro de Ornitologia*. Universidade Federal de Ouro Preto, Ouro Preto, p.8.
- HOFFMANN, D. & RODRIGUES, M. 2006b. Sucesso reprodutivo de *Polystictus superciliaris* (Aves: Tyrannidae) em área de canga no sudeste do Brasil. In: Ribon, R. (Ed.), *Resumos do XIV Congresso Brasileiro de Ornitologia*. Universidade Federal de Ouro Preto, Ouro Preto, p.9.
- HOFFMANN, D. & RODRIGUES, M. 2006c. Território e uso de habitat por *Polystictus superciliaris* (Aves: Tyrannidae) em área de canga no sudeste do Brasil. In: Ribon, R. (Ed.), *Resumos do XIV Congresso Brasileiro de Ornitologia*. Universidade Federal de Ouro Preto, Ouro Preto, p.19.
- HOFFMANN, D. 2006. *Comportamento de forrageamento, dieta, área de vida, biologia reprodutiva e sucesso reprodutivo de Polystictus superciliaris Wied, 1831 [sic] (Aves: Tyrannidae), no sudeste do Brasil*. (M.Sc. Dissertation). Universidade Federal de Minas Gerais, Belo Horizonte.
- HOFFMANN, D.; VASCONCELOS, M.F.; LOPES, L.E. & RODRIGUES, M. 2007. Comportamento de forrageamento e dieta de *Polystictus superciliaris* (Aves, Tyrannidae) no sudeste do Brasil. *Iheringia, Série Zoologia*, 97:296-300.
- HOLT, E.G. 1928. An ornithological survey of the Serra do Itatiaia, Brazil. *Bulletin of the American Museum of Natural History*, 57:251-326.
- HOOGHIEMSTRA, H. & CLEEF, A.M. 1995. Pleistocene climatic change and environmental and generic dynamics in the North Andean montane forest and páramo. In: Churchill, S.P.; Balslev, H.; Forero, E. & Luteyn, E. (Eds.), *Biodiversity and conservation of Neotropical montane forests*. New York Botanical Garden, New York, p.35-49.
- IHERING, H. VON. 1900. Aves observadas em Cantagallo e Nova Friburgo. *Revista do Museu Paulista*, 4:149-164.
- JACOBI, C.M.; CARMO, F.F.; VINCENT, R.C. & STEHMANN, J.R. 2007. Plant communities on ironstone outcrops: a diverse and endangered Brazilian ecosystem. *Biodiversity and Conservation*, 16:2185-2200.
- LEDRU, M.-P.; SALGADO-LABORIAU, M.L. & LORSCHITTEIR, M.L. 1998. Vegetation dynamics in southern and central Brazil during the last 10,000 yr B.P. *Review of Palaeobotany and Palynology*, 99:131-142.
- LENCIONI-NETO, F. 1996. A new subspecies of *Knipolegus* from estado da Bahia, Brasil. *Revista Brasileira de Biologia*, 56:197-201.
- LEONI, L.S. 1997. Táxons endêmicos no Parque Nacional do Caparaó – Minas Gerais. In: Salino, A. (Ed.), *Resumos do 19º Encontro Regional de Botânicos*. Universidade Federal de Minas Gerais, Belo Horizonte, p.43.
- LOPES, L.E. 2008. The range of the Curl-crested Jay: lessons for evaluating bird endemism in the South American Cerrado. *Diversity and Distributions*, 14:561-568.
- LÜDERWALDT, H. 1909. Beitrag zur Ornithologie des Campo Itatiaia. *Zoologischen Jahrbücher*, 27:329-360.
- MACARTHUR, R.H. & WILSON, E.O. 1967. *The theory of island biogeography*. Princeton University Press, Princeton.
- MACHADO, A.B.M.; MARTINS, C.S. & DRUMMOND, G.M. 2005. *Lista da fauna brasileira ameaçada de extinção: incluindo as listas de espécies quase ameaçadas e deficientes em dados*. Fundação Biodiversitas, Belo Horizonte.
- MACHADO, C.G. 2003. Os atributos florais das espécies de plantas visitadas por beija-flores em uma área de campo rupestre da Chapada Diamantina, Bahia. In: Machado, C.G. (Ed.), *Resumos do XI Congresso Brasileiro de Ornitologia*. Universidade Estadual de Feira de Santana, Feira de Santana, p.702.
- MACHADO, C.G. 2005. Aves. In: Juncá, F.A.; Funch, L.S. & Rocha, W. (Eds.), *Biodiversidade e conservação da Chapada Diamantina*. MMA, Brasília, p.357-375.
- MACHADO, C.G.; COELHO, A.G. & SANTANA, C.S. 2003a. Agonismos entre beija-flores em uma área de campo rupestre da Chapada Diamantina, BA. In: Machado, C.G. (Ed.), *Resumos do XI Congresso Brasileiro de Ornitologia*. Universidade Estadual de Feira de Santana, Feira de Santana, p.19.
- MACHADO, C.G.; COELHO, A.G.; SANTANA, C.S. & RODRIGUES, M. 2007a. Beija-flores e seus recursos florais em uma área de campo rupestre da Chapada Diamantina, Bahia. *Revista Brasileira de Ornitologia*, 15:267-279.
- MACHADO, C.G.; MOREIRA, T.A.; NUNES, C.E.C. & ROMÃO, C.O. 2003b. Use of *Micranthocereus purpureus* (Guerke) F. Ritter, 1968 (Cactaceae) hairs in nests of *Augastes lumachellus* Lesson, 1839 [sic] (Trochilidae, Aves). *Sitientibus, Série Ciências Biológicas*, 3:131-132.
- MACHADO, C.G.; NUNES, C.E.C.; GOMES, M.C.B.; COELHO, A.G.; SANTANA, C.S.; MOURA, A.D.C.; ALMEIDA, J.J.; FAUSTINO, T.C. & BASTOS, S.S. 2007b. As espécies de plantas utilizadas por *Phaethornis pretrei* em três tipos de ambientes da Chapada Diamantina. In: Fontana, C.S. (Ed.), *Resumos do XV Congresso Brasileiro de Ornitologia*. Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, p.245.
- MACHADO, R.B.; RIGUEIRA, S.E. & LINS, L.V. 1998. Expansão geográfica do canário-rabudo (*Embernagra longicauda* – Aves: Emberizidae) em Minas Gerais. *Ararajuba*, 6:42-45.
- MALLET-RODRIGUES, F.; PARRINI, R. & PACHECO, J.F. 2007. Birds of the Serra dos Órgãos, state of Rio de Janeiro, southeastern Brazil: a review. *Revista Brasileira de Ornitologia*, 15:5-35.
- MARINI, M.Á. & LOPES, L.E. 2005. Novo limite sul na distribuição geográfica de *Sakesphorus cristatus* (Thamnophilidae). *Ararajuba*, 13:105-106.
- MARTINELLI, G. & ORLEANS E BRAGANÇA, J. 1996. *Campos de altitude*. Editora Index, Rio de Janeiro.
- MATTOS, G.T. & SICK, H. 1985. Sobre a distribuição e a ecologia de duas espécies crípticas: *Embernagra longicauda* Strickland, 1844, e *Embernagra platensis* (Gmelin, 1789). *Emberizidae, Aves. Revista Brasileira de Biologia*, 45:201-206.
- MEGURO, M.; PIRANI, J.R.; GIULIETTI, A.M. & MELLO-SILVA, R. 1994. Phytophysognomy and composition of the vegetation of Serra do Ambrósio, Minas Gerais, Brazil. *Revista Brasileira de Botânica*, 17:149-166.
- MELO-JÚNIOR, T.A.; MENDES, L.G.M. & COELHO, M.M. 1998. Range extension for Itatiaia Spinetail *Oreophylax moreirae* with comments on its distribution. *Cotinga*, 10:68-70.
- MELO-JÚNIOR, T.A.; VASCONCELOS, M.F.; FERNANDES, G.W. & MARINI, M.Á. 2001. Bird species distribution and conservation in Serra do Cipó, Minas Gerais, Brazil. *Bird Conservation International*, 11:189-204.
- MENEZES, N.L. & GIULIETTI, A.M. 1986. Campos rupestres, paraíso botânico na Serra do Cipó. *Ciência Hoje*, 5:38-44.
- MENEZES, N.L. & GIULIETTI, A.M. 2000. Campos rupestres. In: Mendonça, M.P. & Lins, L.V. (Eds.), *Lista vermelha das espécies ameaçadas de extinção da flora de Minas Gerais*. Fundação

- Biodiversitas & Fundação Zoobotânica de Belo Horizonte, Belo Horizonte, p.65-73.
- MIRANDA-RIBEIRO, A. 1906. Vertebrados do Itatiaia (Peixes, Serpentes, Saurios, Aves e Mamíferos). *Archivos do Museu Nacional*, 13:163-190.
- MIRANDA-RIBEIRO, A. 1923. Nota crítica sobre a ornis do Itatiaia. *Archivos do Museu Nacional*, 24:238-255.
- MIRANDA-RIBEIRO, A. 1930. Notas ornithológicas X – ainda *Scytalopus speluncae*. *Boletim do Museu Nacional*, 6:11-16.
- NAUMBURG, E.M.B. 1937. Studies of birds from eastern Brazil and Paraguay, based on a collection made by Emil Kaempfer. *Bulletin of the American Museum of Natural History*, 74:139-205.
- NAUMBURG, E.M.B. 1939. Studies of birds from eastern Brazil and Paraguay, based on a collection made by Emil Kaempfer. *Bulletin of the American Museum of Natural History*, 76:231-276.
- NORES, M. 1995. Insular biogeography of birds on mountain-tops in north western Argentina. *Journal of Biogeography*, 22:61-70.
- O'BRIEN, C.E. 1968. Rediscovery of *Embernagra longicauda* Strickland. *The Auk*, 85:323.
- OLMOS, F. 2007. *Oreophylax moreirae* (Furnariidae) na Serra da Mantiqueira entre Minas Gerais e São Paulo. *Atualidades Ornitológicas*, 137:22-23.
- PACHECO, J.F. & BAUER, C. 1998. Limites setentrionais inéditos e documentados de aves da região da Mata Atlântica no Espírito Santo. *Atualidades Ornitológicas*, 86:4.
- PACHECO, J.F. 2000. Sútis conexões entre W. E. Leach e a ornitologia Brasileira. *Atualidades Ornitológicas*, 94:5.
- PACHECO, J.F.; PARRINI, R.; LOPES, L.E. & VASCONCELOS, M.F. 2008. A avifauna do Parque Estadual do Ibitipoca e áreas adjacentes, Minas Gerais, Brasil, com uma revisão crítica dos registros prévios e comentários sobre biogeografia e conservação. *Cotinga*, 30:16-32.
- PARRINI, R.; RAPOSO, M.A.; PACHECO, J.F.; CARVALHÃES, A.M.P.; MELO-JÚNIOR, T.A.; FONSECA, P.S.M. & MINNS, J.C. 1999. Birds of the Chapada Diamantina, Bahia, Brazil. *Cotinga*, 11:86-95.
- PEARMAN, M. 1990. Behaviour and vocalizations of an undescribed *Canastero Asthenes* sp. from Brazil. *Bulletin of the British Ornithologists' Club*, 110:145-153.
- PEIXOTO-VELHO, P.P. 1923a. Avifauna da Serra do Itatiaia. *Archivos do Museu Nacional*, 24:259-264.
- PEIXOTO-VELHO, P.P. 1923b. Breve notícia sobre a ornis do Caparaó. *Boletim do Museu Nacional*, 1:23-26.
- PETRI, S. & FÚLFARO, V.J. 1986. *Geologia do Brasil (Fanerozóico)*. Editora da Universidade de São Paulo, São Paulo.
- PINESCHI, R.B. 1990. Aves como dispersores de sete espécies de *Rapanea* (Myrsinaceae) no Maciço do Itatiaia, estados do Rio de Janeiro e Minas Gerais. *Aranajuba*, 1:73-78.
- PINTO, O.M.O. 1951. Aves do Itatiaia – lista remissiva e novas achegas à avifauna da região. *Papéis Avulsos do Departamento de Zoologia, São Paulo*, 10:155-208.
- PINTO, O.M.O. 1952. Súmula histórica e sistemática da ornitologia de Minas Gerais. *Arquivos de Zoologia, São Paulo*, 8:1-51.
- PINTO, O.M.O. 1954. Aves do Itatiaia – lista remissiva e novas achegas à avifauna da região. *Boletim do Parque Nacional do Itatiaia*, 3:1-87.
- PINTO, O.M.O. 1978. *Novo catálogo das aves do Brasil*. Empresa Gráfica da Revista dos Tribunais, São Paulo. Primeira parte.
- PIRATELLI, A.J. 1997. Comportamento alimentar de beija-flores em duas espécies de *Hippeastrum* Herb. (Amaryllidaceae). *Revista Brasileira de Biologia*, 57:261-273.
- RAPOSO, M.A.; STOPIGLIA, R.; LOSKOT, V. & KIRWAN, G.M. 2006. The correct use of the name *Scytalopus speluncae* (Ménétriéri, 1835), and the description of a new species of Brazilian tapaculo (Aves: Passeriformes: Rhinocryptidae). *Zootaxa*, 1271:37-56.
- REMSEN, J.V. 1994. Use and misuse of bird lists in community ecology and conservation. *The Auk*, 111:225-227.
- RIBEIRO, J.F. & WALTER, B.M.T. 1998. Fitofisionomias do bioma Cerrado. In: Sano, S.M. & Almeida, S.P. (Eds.), *Cerrado: ambiente e flora*. EMBRAPA – CPAC, Planaltina.
- RIBEIRO, K.T.; MEDINA, B.M.O. & SCARANO, F.R. 2007. Species composition and biogeographic relations of the rock outcrop flora on the high plateau of Itatiaia, SE-Brazil. *Revista Brasileira de Botânica*, 30:623-639.
- RIBEIRO, R.C.C. 1997. Ocorrência de águia-cinzenta (*Harpheiaetus [sic] coronatus*) no município de Itambé do Mato Dentro, MG. *Atualidades Ornitológicas*, 78:14.
- RIBON, R.; COELHO, F.M.; PIZETTA, G.T.; ALMEIDA, L.B.; OLIVEIRA, A.; RESCK, R.P.; VALADARES, C.F.; CORRAL, L.; AMBONI, M.; SOUZA, R.A. & CARVALHO, L.F. 2006. Vocalizações, territórios e densidade de *Anthus hellmayri* (Motacillidae: Passeriformes) no Parque Nacional da Serra do Cipó, Brasil. In: Ribon, R. (Ed.), *Resumos do XIV Congresso Brasileiro de Ornitologia*. Universidade Federal de Ouro Preto, Ouro Preto, p.46-47.
- RIDGELY, R.S. & TUDOR, G. 1989. *The birds of South America*. University of Texas Press, Austin, v.1.
- RIDGELY, R.S. & TUDOR, G. 1994. *The birds of South America*. University of Texas Press, Austin, v.2.
- RIZZINI, C.T. 1997. *Tratado de fitogeografia do Brasil: aspectos ecológicos, sociológicos e florísticos*. Âmbito Cultural Edições Ltda, Rio de Janeiro.
- ROMÃO, C.O.; GIMENES, M. & MACHADO, C.G. 2001. Interações entre *Augastes lumachellus* e seus recursos florais no Morro do Pai Inácio, Chapada Diamantina, Bahia. In: Straube, F.C. (Ed.), *Ornitologia sem fronteiras (incluindo os resumos do IX Congresso Brasileiro de Ornitologia)*. Pontifícia Universidade Católica do Paraná, Curitiba, p.336-337.
- RUSCHI, A. 1951. Trochilídeos do Museu Nacional. *Boletim do Museu de Biologia Prof. Mello Leitão*, Série Biologia, 10:1-111.
- RUSCHI, A. 1962a. Algumas observações sobre *Augastes lumachellus* (Lesson) e *Augastes scutatus* (Temminck). *Boletim do Museu de Biologia Prof. Mello Leitão*, Série Biologia, 31:1-24.
- RUSCHI, A. 1962b. Um novo representante de *Colibri* (Trochilidae Aves) da região de Andaraí no estado da Bahia. *Boletim do Museu de Biologia Prof. Mello Leitão*, Série Biologia, 32:1-7.
- RUSCHI, A. 1963a. Notes on Trochilidae: the genus *Augastes*. *Proceedings of the XIII International Ornithological Congress*, 1:141-146.
- RUSCHI, A. 1963b. A atual distribuição geográfica das espécies e sub-espécies do gênero *Augastes*, com a descrição de uma nova sub-espécie: *Augastes scutatus soaresi* Ruschi e a chave artificial e analítica para o reconhecimento das mesmas. (Trochilidae: Aves). *Boletim do Museu de Biologia Prof. Mello Leitão*, Série Divulgação, 4:1-4.
- RUSCHI, A. 1963c. Um novo representante de *Campylopterus*, da região de Diamantina, no estado de Minas Gerais (Trochilidae: Aves). *Boletim do Museu de Biologia Prof. Mello Leitão*, Série Biologia, 39:1-9.
- RUSCHI, A. 1964. O genero *Campylopterus* e as espécies representadas no Brasil. A sua atual distribuição geográfica, com um novo representante para o Brasil (Trochilidae: Aves). *Boletim do Museu de Biologia Prof. Mello Leitão*, Série Biologia, 40:1-5.
- RUSCHI, A. 1975. *Phaethornis pretrei schwarti* n.s.sp. *Boletim do Museu de Biologia Prof. Mello Leitão*, Série Zoologia, 82:1-4.
- RUSCHI, A. 1978. Mamíferos e aves do Parque Nacional do Caparaó. *Boletim do Museu de Biologia Prof. Mello Leitão*, Série Zoologia, 95:1-28.

- RUSCHI, A. 1982a. *Aves do Brasil: beija-flores*. Expressão e Cultura, Rio de Janeiro, v.5.
- RUSCHI, A. 1982b. Mamíferos e aves observadas na Reserva Biológica de Pedra Azul no estado do Espírito Santo. *Boletim do Museu de Biologia Prof. Mello Leitão*, Série Zoológica, 103:1-15.
- SAFFORD, H.D. 1999a. Brazilian páramos I. An introduction to the physical environment and vegetation of the campos de altitude. *Journal of Biogeography*, 26:693-712.
- SAFFORD, H.D. 1999b. Brazilian páramos II. Macro- and mesoclimate of the campos de altitude and affinities with high mountain climates of the tropical Andes and Costa Rica. *Journal of Biogeography*, 26:713-737.
- SAFFORD, H.D. 2007. Brazilian páramos IV. Phytogeography of the campos de altitude. *Journal of Biogeography*, 34:1701-1722.
- SANTANA, C.S. & MACHADO, C.G. 2006. Fenologia de floração e os visitantes florais de *Orthophytum albopictum* Philcox (Bromeliaceae) em uma área de campo rupestre da Chapada Diamantina, em Mucugê, BA. In: *Resumos do 57 Congresso Brasileiro de Botânica*, Gramado.
- SANTOS, A.S.R. 2000. Primeiro registro documentado da garrincha-chorona, *Schizoeaca moreirae*, para o estado de São Paulo. (Passeriformes: Furnariidae). *Boletim CEO*, 14:51-52.
- SAZIMA, M. & SAZIMA, I. 1990. Hummingbird pollination in two species of *Vellozia* (Liliiflorae: Velloziaceae) in southeastern Brazil. *Botanica Acta*, 103:83-86.
- SAZIMA, M. 1977. Hummingbird pollination of *Barbacenia flava* (Velloziaceae) in the Serra do Cipó, Minas Gerais, Brazil. *Flora*, 166:239-247.
- SCHUBART, O.; AGUIRRE, Á.C. & SICK, H. 1965. Contribuição para o conhecimento da alimentação das aves brasileiras. *Arquivos de Zoologia, São Paulo*, 12:95-249.
- SICK, H. 1959. O redescobrimto no Brasil do bacurau *Caprimulgus longirostris* Bonaparte (Caprimulgidae, Aves). *Boletim do Museu Nacional*, Nova Série Zoológica, 204:1-15.
- SICK, H. 1963. O bacurau *Caprimulgus longirostris* Bon. e outras aves noturnas do estado da Guanabara (GB). *Vellozia*, 1:107-116.
- SICK, H. 1970. Der Strohschwanz, *Oreophylax moreirae*, andiner Furnariidae in südostbrasilien. *Bonner Zoologische Beitrage*, 21:251-268.
- SICK, H. 1985. Observations on the Andean-Patagonian component of southeastern Brazil's avifauna. *Ornithological Monographs*, 36:233-237.
- SICK, H. 1997. *Ornitologia Brasileira*. Nova Fronteira, Rio de Janeiro.
- SILVA, J.M.C. & BATES, J.M. 2002. Biogeographic patterns and conservation in the South American Cerrado: a tropical savanna hotspot. *BioScience*, 52:225-233.
- SILVA, J.M.C. & SANTOS, M.P.D. 2005. A importância relativa dos processos biogeográficos na formação da avifauna do Cerrado e de outros biomas brasileiros. In: Scariot, A.; Sousa Filho, J.C. & Felfili, J.M. (Eds.), *Cerrado: ecologia, biodiversidade e conservação*. Ministério do Meio Ambiente, Brasília, p.220-233.
- SILVA, J.M.C. & STOTZ, D.F. 1992. Geographic variation in the Sharp-billed Treehunter *Heliobletus contaminatus*. *Bulletin of the British Ornithologists' Club*, 112:98-101.
- SILVA, J.M.C. & STRAUBE, F.C. 1996. Systematics and biogeography of Scaled Woodcreepers (Aves: Dendrocolaptidae). *Studies on Neotropical Fauna and Environment*, 31:3-10.
- SILVA, J.M.C. 1995. Biogeographic analysis of the South American Cerrado avifauna. *Steenstrupia*, 21:49-67.
- SILVA, J.M.C. 1997. Endemic bird species and conservation in the Cerrado region, South America. *Biodiversity and Conservation*, 6:435-450.
- SILVA, J.M.C. 1998. Integrating biogeography and conservation: an example with birds and plants of the Cerrado region. *Anais da Academia Brasileira de Ciências*, 70:881-888.
- SILVEIRA, F.A. & CURE, J.R. 1993. High altitude bee fauna of Southeastern Brazil: implications for biogeographic patterns (Hymenoptera: Apoidea). *Studies on Neotropical Fauna and Environment*, 28:47-55.
- SILVEIRA, L.F. 1998. The birds of Serra da Canastra National Park and adjacent areas, Minas Gerais, Brazil. *Cotinga*, 10:55-63.
- SIMON, J.E.; RIBON, R.; MATTOS, G.T. & ABREU, C.R.M. 1999. A avifauna do Parque Estadual da Serra do Brigadeiro, Minas Gerais. *Revista Árvore*, 23:33-48.
- SIMPSON, B.B. 1979. Quaternary biogeography of the high montane regions of South America. In: Duellman, W.E. (Ed.), *The South American herpetofauna: its origin, evolution, and dispersal*. University of Kansas, Lawrence, p.157-188.
- SIMPSON-VUILLEUMIER, B. 1971. Pleistocene changes in the fauna and flora of South America. *Science*, 173:771-780.
- SNETHLAGE, E. 1930. Fam. Pteroptochidae. *Boletim do Museu Nacional*, 6:9-10.
- SPIX, J.B. 1825. *Avium species novae, quas in itinere per Brasiliam annis 1817-20 collegit et descripsi*. Hubschmann, München.
- STATTERSFIELD, A.J.; CROSBY, M.J.; LONG, A.J. & WEGE, D.C. 1998. *Endemic bird areas of the world: priorities for biodiversity conservation*. BirdLife International, Cambridge, BirdLife Conservation Series nº 7.
- STEYERMARK, J.A. 1986. Speciation and endemism in the flora of the Venezuelan Tepuis. In: Vuilleumier, F. & Monasterio, M. (Eds.), *High altitude tropical biogeography*. Oxford University Press, New York, p.317-373.
- STILES, F.G. 1999. Brown Violet-ear *Colibri delphinae*. In: del Hoyo, J.; Elliott, A. & Sargatal, J. (Eds.), *Handbook of the birds of the world*. Lynx Edicions, Barcelona, v.5: Barn-owls to hummingbirds, p.557.
- STOTZ, D.F.; FITZPATRICK, J.W.; PARKER III, T.A. & MOSKOVITS, D.K. 1996. *Neotropical birds: ecology and conservation*. University of Chicago Press, Chicago.
- STUDER, A. & TEIXEIRA, D.M. 1993. Notas sobre a biologia reprodutiva de *Asthenes luizae* Viellard, 1990 (Aves, Furnariidae). In: Cirne, M.P. (Ed.), *Resumos do III Congresso Brasileiro de Ornitologia*. Editora da Universidade Católica de Pelotas, Pelotas, p.44.
- ULE, E. 1895. Relatório de uma excursão botânica feita na Serra do Itatiaia. *Arquivos do Museu Nacional*, 9:185-223.
- VAN DER HAMMEN, T. 1974. The Pleistocene changes of vegetation and climate in tropical South America. *Journal of Biogeography*, 1:3-26.
- VASCONCELOS, M.F. & BRANDT, L.F.S. 1998. Distribuição altitudinal e por habitats da avifauna da Serra do Curral, Belo Horizonte, MG. In: Alves, M.A.S. (Ed.), *Resumos do VII Congresso Brasileiro de Ornitologia*. Universidade do Estado do Rio de Janeiro, Rio de Janeiro, p.142.
- VASCONCELOS, M.F. & D'ANGELO-NETO, S. 2007. Padrões de distribuição e conservação da avifauna na região central da Cadeia do Espinhaço e áreas adjacentes, Minas Gerais, Brasil. *Cotinga*, 28:27-44.
- VASCONCELOS, M.F. & ENDRIGO, E. 2008. O canário-rasteiro (*Sicalis citrina*) nas serras de Minas Gerais: notas adicionais. *Atualidades Ornitológicas*, 141:14-15.
- VASCONCELOS, M.F. & FERREIRA, J.C. 2001. Sazonalidade e reprodução do andorinhão-de-coleira-falha (*Streptoprocne biscutata*) no Pico do Inficionado, Serra do Caraça, Minas Gerais, Brasil. *Tangara*, 1:74-84.
- VASCONCELOS, M.F. & LOMBARDI, J.A. 1996. Primeira descrição do ninho e do ovo de *Polystictus superciliaris* (Passeriformes:



- Tyrannidae) ocorrente na Serra do Curral, Minas Gerais. *Ararajuba*, 4:114-116.
- VASCONCELOS, M.F. & LOMBARDI, J.A. 1999. Padrão sazonal na ocorrência de seis espécies de beija-flores (Apodiformes: Trochilidae) em uma localidade de campo rupestre na Serra do Curral, Minas Gerais. *Ararajuba*, 7:71-79.
- VASCONCELOS, M.F. & LOMBARDI, J.A. 2000. Espécies vegetais visitadas por beija-flores durante o meio do verão no Parque Estadual da Pedra Azul, Espírito Santo. *Melospittacus*, 3:36-41.
- VASCONCELOS, M.F. & LOMBARDI, J.A. 2001. Hummingbirds and their flowers in the campos rupestres of southern Espinhaço Range, Brazil. *Melospittacus*, 4:3-30.
- VASCONCELOS, M.F. & MELO-JÚNIOR, T.A. 2001. An ornithological survey of Serra do Caraça, Minas Gerais, Brazil. *Cotinga*, 15:21-31.
- VASCONCELOS, M.F. & SILVA, J.A. 2003. Descrição do jovem de Tibirro-rupestre *Embernagra longicauda*. *Cotinga*, 20:21-23.
- VASCONCELOS, M.F. 1997. Nidificações do urubu-de-cabeça-preta (*Coragyps atratus*) em um fosso profundo como uma provável defesa contra incêndios. *Bios*, 5:61-64.
- VASCONCELOS, M.F. 1999a. Contribuição ao conhecimento ornitológico do Pico do Papagaio, município de Aiuruoca, Minas Gerais. *Atualidades Ornitológicas*, 90:10-11.
- VASCONCELOS, M.F. 1999b. Natural history notes and conservation of two species endemic to the Espinhaço Range, Brazil: Hyacinth Visorbearer *Augastes scutatus* and Grey-backed Tachuri *Polystictus superciliiaris*. *Cotinga*, 11:75-78.
- VASCONCELOS, M.F. 2000. Reserva do Caraça: história, vegetação e fauna. *Aves*, 1:3-7.
- VASCONCELOS, M.F. 2001a. Adições à avifauna da Serra do Caraça, Minas Gerais. *Atualidades Ornitológicas*, 104:3-4.
- VASCONCELOS, M.F. 2001b. Pale-throated Serra-finch *Embernagra longicauda*. *Cotinga*, 16:110-112.
- VASCONCELOS, M.F. 2002. O João-cipó (*Asthenes luizae*) no Parque Estadual do Pico do Itambé, Minas Gerais, Brasil. *Atualidades Ornitológicas*, 107:10.
- VASCONCELOS, M.F. 2003. A avifauna dos campos de altitude da Serra do Caparaó, estados de Minas Gerais e Espírito Santo, Brasil. *Cotinga*, 19:40-48.
- VASCONCELOS, M.F. 2007. Aves observadas no Parque Paredão da Serra do Curral, Belo Horizonte, Minas Gerais, Brasil. *Atualidades Ornitológicas*, 136:6-11.
- VASCONCELOS, M.F. 2008a. Aves registradas na Serra do Papagaio, município de Aiuruoca, Minas Gerais. *Atualidades Ornitológicas*, 142:6-7.
- VASCONCELOS, M.F. 2008b. Mountaintop endemism in eastern Brazil: why some bird species from campos rupestres of the Espinhaço Range are not endemic to the Cerrado region? *Revista Brasileira de Ornitologia*, 16:348-362.
- VASCONCELOS, M.F.; D'ANGELO NETO, S. & FIELDS, J. 2008b. Redescription of Cipó Canastero *Asthenes luizae*, with notes on its systematic relationships. *Bulletin of the British Ornithologists' Club*, 128:179-186.
- VASCONCELOS, M.F.; D'ANGELO NETO, S.; KIRWAN, G.M.; BORNSCHEIN, M.R.; DINIZ, M.G. & SILVA, J.F. 2006. Important ornithological records from Minas Gerais state, Brazil. *Bulletin of the British Ornithologists' Club*, 126:212-238.
- VASCONCELOS, M.F.; D'ANGELO-NETO, S. & RODRIGUES, M. 2002a. A range extension for the Cipó Canastero *Asthenes luizae* and the consequences for its conservation status. *Bulletin of the British Ornithologists' Club*, 122:7-10.
- VASCONCELOS, M.F.; FIGUEREDO, C.C. & OLIVEIRA, R.S. 1998. Padrão temporal de vocalização do bacurau-da-telha *Caprimulgus longirostris* (Aves, Caprimulgidae) ao longo de quatro noites na Serra do Curral, Minas Gerais, Brasil. *Boletim do Museu de Biologia Mello Leitão*, Nova Série, 9:13-17.
- VASCONCELOS, M.F.; FIGUEREDO, C.C. & OLIVEIRA, R.S. 1999a. Táticas de forrageamento do bacurau-da-telha *Caprimulgus longirostris* (Aves, Caprimulgidae) na Serra do Curral, Minas Gerais, Brasil. *Boletim do Museu de Biologia Mello Leitão*, Nova Série, 10:33-38.
- VASCONCELOS, M.F.; LOMBARDI, V.T. & D'ANGELO NETO, S. 2007a. Notas sobre o canário-rasteiro (*Sicalis citrina*) nas serras de Minas Gerais, Brasil. *Atualidades Ornitológicas*, 140:6-7.
- VASCONCELOS, M.F.; LOPES, L.E. & HOFFMANN, D. 2007b. Dieta e comportamento de forrageamento de *Oreophylax moreirae* (Aves: Furnariidae) na Serra do Caraça, Minas Gerais, Brasil. *Revista Brasileira de Ornitologia*, 15:439-442.
- VASCONCELOS, M.F.; MALDONADO-COELHO, M. & BUZZETTI, D.R.C. 2003. Range extensions for the Gray-backed Tachuri (*Polystictus superciliiaris*) and the Pale-throated Serra-finch (*Embernagra longicauda*) with a revision on their geographic distribution. *Ornitologia Neotropical*, 14:477-489.
- VASCONCELOS, M.F.; MALDONADO-COELHO, M. & DURÃES, R. 1999b. Notas sobre algumas espécies de aves ameaçadas e pouco conhecidas da porção meridional da Cadeia do Espinhaço, Minas Gerais. *Melospittacus*, 2:44-50.
- VASCONCELOS, M.F.; MAURÍCIO, G.N.; KIRWAN, G.M. & SILVEIRA, L.F. 2008a. Range extension for Marsh Tapaculo *Scytalopus iraiensis* to the highlands of Minas Gerais, Brazil, with an overview of the species' distribution. *Bulletin of the British Ornithologists' Club*, 128:101-106.
- VASCONCELOS, M.F.; SALINO, A. & VIEIRA, M.V.O. 2002b. A redescoberta de *Huperzia rubra* (Cham.) Trevisan (Lycopodiaceae) e o seu atual estado de conservação nas altas montanhas do sul da Cadeia do Espinhaço, Minas Gerais. *Unimontes Científica*, 3:45-50.
- VASCONCELOS, M.F.; VASCONCELOS, P.N. & FERNANDES, G.W. 2001. Observations on a nest of Hyacinth Visorbearer *Augastes scutatus*. *Cotinga*, 16:57-61.
- VERSIEUX, L.M. & WENDT, T. 2007. Bromeliaceae diversity and conservation in Minas Gerais state, Brazil. *Biodiversity and Conservation*, 16:2989-3009.
- VIANA, L.R.; FERNANDES, G.W. & SILVA, C.E. 2005. 'Ecological road' threatens endemic Brazilian plant with extinction. *Plant Talk*, 41:15.
- VIELLIARD, J.M.E. 1990a. Estudo bioacústico das aves do Brasil: o gênero *Scytalopus*. *Ararajuba*, 1:5-18.
- VIELLIARD, J.M.E. 1990b. Uma nova espécie de *Asthenes* da Serra do Cipó, Minas Gerais, Brasil. *Ararajuba*, 1:121-122.
- VIELLIARD, J.M.E. 1994. *Catálogo dos troquilídeos do Museu de Biologia Mello Leitão*. Museu de Biologia Mello Leitão, Santa Teresa.
- VIELLIARD, J.M.E. 1996. Areas of differentiation and biogeographic affinities within the avifauna of northeastern Brazil. In: Vielliard, J.M.E.; Silva, M.L. & Silva, W.R. (Eds.), *Anais do V Congresso Brasileiro de Ornitologia*. Universidade Estadual de Campinas, Campinas, p.184-190.
- VINCENT, R.C. 2004. *Florística, fitossociologia e relações entre a vegetação e o solo em áreas de campos ferruginosos no Quadrilátero Ferrífero, Minas Gerais*. (Ph.D. Dissertation). Universidade de São Paulo, São Paulo.
- VUILLEUMIER, F. 1970. Insular biogeography in continental regions. I. The northern Andes of South America. *The American Naturalist*, 104:373-388.
- VUILLEUMIER, F. 1986. Origins of the tropical avifaunas of the high Andes. In: Vuilleumier, F. & Monasterio, M. (Eds.), *High altitude tropical biogeography*. Oxford University Press, New York, p.586-622.

- WIED, M. PRINZ ZU. 1831. *Beiträge zur Naturgeschichte von Brasilien, Vögel*. Landes Industrie Comptoirs, Weimar.
- WILLIS, E.O. & ONIKI, Y. 1991. Avifaunal transects across the open zones of northern Minas Gerais, Brazil. *Ararajuba*, 2:41-58.
- WILLIS, E.O. 1992. Zoogeographical origins of eastern Brazilian birds. *Ornitologia Neotropical*, 3:1-15.
- WILLIS, E.O. 2002. Birds at *Eucalyptus* and other flowers in southern Brazil: a review. *Ararajuba*, 10:43-66.
- ZORZIN, G.; CARVALHO, C.E.A.; CARVALHO-FILHO, E.P.M. & CANUTO, M. 2006. Novos registros de Falconiformes raros e ameaçados para o estado de Minas Gerais. *Revista Brasileira de Ornitologia*, 14:417-421.

Recebido em: 11.07.2008

Aceito em: 18.11.2009

Impresso em: 31.03.2010

## APPENDIX 1

List of sources used in the revision of the open-habitat avifauna of southeastern Brazilian mountaintops.

Number	Source	Number	Source
1	Spix (1825)	51	Lencioni-Neto (1996)
2	Wied (1831)	52	Martinelli & Orleans e Bragança (1996)
3	Ule (1895)	53	Vasconcelos & Lombardi (1996)
4	Hemmendorff & Moreira (1903)	54	Andrade (1997a)
5	Miranda-Ribeiro (1906)	55	Andrade (1997b)
6	Gounelle (1909)	56	Piratelli (1997)
7	Lüderwaldt (1909)	57	Ribeiro (1997)
8	Miranda-Ribeiro (1923)	58	Sick (1997)
9	Peixoto-Velho (1923a)	59	Vasconcelos (1997)
10	Peixoto-Velho (1923b)	60	Andrade (1998)
11	Holt (1928)	61	Andrade <i>et al.</i> (1998)
12	Miranda-Ribeiro (1930)	62	Cordeiro <i>et al.</i> (1998)
13	Snethlage (1930)	63	Machado <i>et al.</i> (1998)
14	Naumburg (1937)	64	Melo-Júnior <i>et al.</i> (1998)
15	Naumburg (1939)	65	Pacheco & Bauer (1998)
16	Pinto (1951)	66	Vasconcelos & Brandt (1998)
17	Ruschi (1951)	67	Vasconcelos <i>et al.</i> (1998)
18	Pinto (1952)	68	Almeida & Raposo (1999)
19	Pinto (1954)	69	Bauer (1999)
20	Sick (1959)	70	Funch (1999)
21	Ruschi (1962a)	71	Parrini <i>et al.</i> (1999)
22	Ruschi (1962b)	72	Simon <i>et al.</i> (1999)
23	Ruschi (1963a)	73	Vasconcelos (1999a)
24	Ruschi (1963b)	74	Vasconcelos (1999b)
25	Ruschi (1963c)	75	Vasconcelos & Lombardi (1999)
26	Sick (1963)	76	Vasconcelos <i>et al.</i> (1999a)
27	Ruschi (1964)	77	Vasconcelos <i>et al.</i> (1999b)
28	Schubart <i>et al.</i> (1965)	78	Pacheco (2000)
29	Grantsau (1967)	79	Santos (2000)
30	Grantsau (1968)	80	Vasconcelos & Lombardi (2000)
31	O'Brien (1968)	81	Vasconcelos (2000)
32	Sick (1970)	82	Carvalhoes (2001)
33	Ruschi (1975)	83	Gonzaga & Castiglioni (2001)
34	Sazima (1977)	84	Melo-Júnior <i>et al.</i> (2001)
35	Carnevali (1980)	85	Romão <i>et al.</i> (2001)
36	Carnevali (1982)	86	Vasconcelos (2001a)
37	Ruschi (1982a)	87	Vasconcelos (2001b)
38	Mattos & Sick (1985)	88	Vasconcelos & Ferreira (2001)
39	Dorst & Vuilleumier (1986)	89	Vasconcelos & Lombardi (2001)
40	Grantsau (1988)	90	Vasconcelos & Melo-Júnior (2001)
41	Ridgely & Tudor (1989)	91	Vasconcelos <i>et al.</i> (2001)
42	Pearman (1990)	92	Brammer (2002)
43	Pineschi (1990)	93	Vasconcelos (2002)
44	Sazima & Sazima (1990)	94	Vasconcelos <i>et al.</i> (2002a)
45	Vielliard (1990a)	95	Willis (2002)
46	Vielliard (1990b)	96	Coelho & Machado (2003)
47	Willis & Oniki (1991)	97	Faustino & Machado (2003)
48	Studer & Teixeira (1993)	98	Faustino <i>et al.</i> (2003)
49	Ridgely & Tudor (1994)	99	Machado (2003)
50	Vielliard (1994)	100	Machado <i>et al.</i> (2003a)

Number	Source	Number	Source
101	Machado <i>et al.</i> (2003b)	135	Machado <i>et al.</i> (2007b)
102	Vasconcelos (2003)	136	Mallet-Rodrigues <i>et al.</i> (2007)
103	Vasconcelos & Silva (2003)	137	Olmos (2007)
104	Vasconcelos <i>et al.</i> (2003)	138	Vasconcelos (2007)
105	Hoffmann & Rodrigues (2005)	139	Vasconcelos & D'Angelo-Neto (2007)
106	Machado (2005)	140	Vasconcelos <i>et al.</i> (2007a)
107	Abreu (2006)	141	Vasconcelos <i>et al.</i> (2007b)
108	Alves <i>et al.</i> (2006)	142	Freitas & Rodrigues (2008)
109	Colaço <i>et al.</i> (2006)	143	Vasconcelos (2008a)
110	Costa & Rodrigues (2006a)	144	Vasconcelos & Endrigo (2008)
111	Costa & Rodrigues (2006b)	145	Costa & Rodrigues ( <i>pers. com.</i> )
112	Domingues & Rodrigues (2006)	146	Costa <i>et al.</i> (2008)
113	Faustino & Machado (2006)	147	Freitas <i>et al.</i> (2008)
114	Gomes (2006)	148	Pacheco <i>et al.</i> (2008)
115	Gomes & Guerra (2006)	149	Vasconcelos <i>et al.</i> (2008a)
116	Gomes & Rodrigues (2006a)	150	Vasconcelos <i>et al.</i> (2008b)
117	Gomes & Rodrigues (2006b)	151	AMNH
118	Guerra & Alves (2006)	152	DZUFMG
119	Guerra <i>et al.</i> (2006)	153	MBML
120	Hoffmann (2006)	154	MCN
121	Hoffmann & Rodrigues (2006a)	155	MCP
122	Hoffmann & Rodrigues (2006b)	156	MHNT
123	Hoffmann & Rodrigues (2006c)	157	MNRJ
124	Raposo <i>et al.</i> (2006)	158	MPEG
125	Ribon <i>et al.</i> (2006)	159	MZUSP
126	Santana & Machado (2006)	160	SG
127	Vasconcelos <i>et al.</i> (2006)	161	F.F. Vasconcelos ( <i>pers. com.</i> )
128	Zorzin <i>et al.</i> (2006)	162	F. Mallet-Rodrigues ( <i>pers. com.</i> )
129	Alves, A.C.F. <i>et al.</i> (2007)	163	F.S. Leite ( <i>pers. com.</i> )
130	Assis <i>et al.</i> (2007)	164	G.B. Malacco ( <i>pers. com.</i> )
131	Costa & Rodrigues (2007)	165	J.F. Pacheco ( <i>pers. com.</i> )
132	Gonzaga <i>et al.</i> (2007)	166	R. Parrini ( <i>pers. com.</i> )
133	Hoffmann <i>et al.</i> (2007)	167	Personal observation
134	Machado <i>et al.</i> (2007a)		

APPENDIX 2

Checklist of the avifauna recorded in the *campos rupestres* and *campos de altitude* of southeastern Brazilian mountaintops and patterns of endemism. Numerical codes for localities and sources are presented in Table 1 and Appendix 1, respectively. Endemism: AF = endemic to the Atlantic Forest region; CA = endemic to the Caatinga region; CE = endemic to the Cerrado region; MT = endemic to the open-habitats of southeastern Brazilian mountaintops (*campos rupestres* and *campos de altitude*).

Family/Species	<i>Campos rupestres</i>	<i>Campos de altitude</i>	Sources	Endemism
Tinamidae				
<i>Crypturellus parvirostris</i>	I.5, I.10, I.15, I.18, I.20, I.23, I.24, I.25, I.27, I.32, I.35, I.36, I.37, I.43, I.44		70, 71, 84, 106, 139, 167	
<i>Rhynchotus rufescens</i>	I.10, I.15, I.24, I.25, I.26, I.33, I.35, I.37, I.40, I.44	II.6, II.8	3, 5, 7, 8, 11, 16, 19, 39, 58, 70, 71, 82, 84, 139, 145, 148, 167	
<i>Nothura minor</i>	I.30		1, 18	CE
<i>Nothura maculosa</i>	I.5, I.10, I.25, I.26, I.30, I.32, I.35, I.50	II.6, II.8	5, 8, 11, 16, 18, 19, 54, 84, 106, 139, 145, 148, 152, 167	
Threskiornithidae				
<i>Theristicus caudatus</i>	I.32		93, 167	
Cathartidae				
<i>Cathartes aura</i>	I.10, I.15, I.18, I.20, I.24, I.25, I.26, I.32, I.35, I.37, I.42, I.43, I.44, I.46, I.50, I.51	II.1, II.6, II.8	11, 16, 19, 47, 84, 90, 106, 139, 145, 148, 167	
<i>Coragyps atratus</i>	I.3, I.5, I.10, I.18, I.20, I.24, I.25, I.26, I.27, I.35, I.36, I.37, I.38, I.40, I.42, I.43, I.46, I.47, I.48, I.50, I.51	II.2, II.6, II.7, II.8, II.11, II.12, III.1, III.2	5, 8, 11, 16, 19, 39, 47, 59, 70, 71, 84, 90, 106, 136, 138, 139, 145, 148, 162, 164, 166, 167	
<i>Sarcoramphus papa</i>	I.3, I.27, I.35, I.42, I.46, I.48		70, 71, 84, 90, 145, 167	
Accipitridae				
<i>Elanus leucurus</i>		II.6	148	
<i>Ictinia plumbea</i>	I.5		106	
<i>Geranospiza caerulescens</i>	I.32, I.35		61, 167	
<i>Heterospizias meridionalis</i>	I.35	II.6	145, 148	
<i>Harpyhaliaetus coronatus</i>	I.5, I.35, I.39		57, 71, 128	
<i>Percnophierax leucorrhous</i>		II.8	11, 16, 19	
<i>Rupornis magnirostris</i>	I.5, I.10, I.15, I.20, I.21, I.24, I.25, I.26, I.27, I.32, I.35, I.36, I.38, I.40, I.42, I.43, I.44, I.46, I.47, I.48, I.50	II.6, II.8, II.11	84, 86, 90, 106, 138, 139, 145, 148, 164, 167	
<i>Buteo albicaudatus</i>	I.18, I.26, I.27, I.32, I.35, I.36, I.38, I.40, I.42, I.44, I.46, I.50, I.51	II.1, II.2, II.6, II.8, II.10, II.11, III.1, III.2	47, 52, 84, 90, 136, 137, 138, 139, 145, 148, 162, 164, 167	
<i>Buteo melanoleucus</i>	I.5, I.15, I.20, I.25, I.35, I.38, I.42, I.46	II.6	47, 70, 71, 82, 88, 90, 138, 139, 145, 148, 167	
<i>Buteo albonotatus</i>	I.18, I.35, I.36, I.42		84, 86, 90, 145, 167	
Falconidae				
<i>Caracara plancus</i>	I.3, I.10, I.26, I.27, I.33, I.35, I.36, I.37, I.38, I.40, I.42, I.43, I.44, I.46, I.48, I.50, I.51	II.6, II.7, II.8	70, 71, 82, 84, 90, 106, 138, 139, 143, 145, 148, 164, 167	
<i>Milvago chimachima</i>	I.3, I.5, I.10, I.15, I.24, I.25, I.26, I.27, I.32, I.33, I.35, I.36, I.38, I.40, I.42, I.43, I.50, I.51	II.6, II.8	7, 8, 47, 70, 71, 82, 84, 86, 90, 138, 139, 145, 148, 164, 167	
<i>Herpetotheres cachinnans</i>	I.20, I.35, I.42	II.6, II.7	90, 145, 148, 167	
<i>Falco sparverius</i>	I.3, I.10, I.15, I.25, I.27, I.32, I.35, I.36, I.38, I.42, I.43, I.44, I.49	II.6, II.7, II.8	7, 8, 11, 16, 19, 39, 47, 71, 82, 84, 90, 138, 145, 148, 152, 167	

Family/Species	<i>Campos rupestres</i>	<i>Campos de altitude</i>	Sources	Endemism
<i>Falco femoralis</i>	I.15, I.27, I.35, I.38, I.42	II.6	47, 70, 71, 82, 90, 138, 148, 167	
<i>Falco peregrinus</i>	I.42		86, 167	
Rallidae				
<i>Porzana albicollis</i>	I.35		167	
Cariamidae				
<i>Cariama cristata</i>	I.5, I.10, I.15, I.20, I.24, I.26, I.27, I.32, I.33, I.35, I.36, I.37, I.42, I.43, I.50, I.51	II.1, II.6, II.7, II.8, III.2, III.3	7, 11, 16, 19, 52, 58, 70, 71, 82, 84, 90, 106, 136, 139, 143, 145, 148, 162, 167	
Scolopacidae				
<i>Gallinago paraguayiae</i>	I.15, I.27		70, 71, 167	
<i>Gallinago undulata</i>	I.32, I.35		167	
Columbidae				
<i>Columbina minuta</i>	I.10		71, 82	
<i>Columbina talpacoti</i>	I.35, I.38, I.40	II.1	10, 47, 84, 102, 164, 167	
<i>Columbina squammata</i>	I.32, I.35, I.43		84, 167	
<i>Columbina picui</i>	I.5, I.10, I.23, I.24		106, 139, 167	
<i>Patagioenas picazuro</i>	I.26, I.27, I.35, I.36, I.38, I.40, II.5		84, 138, 139, 145, 164, 167	
<i>Patagioenas cayennensis</i>	I.35, I.42	II.6	84, 90, 145, 148	
<i>Patagioenas plumbea</i>	I.40	II.8	5, 7, 8, 11, 16, 19, 43, 164	
<i>Zenaida auriculata</i>	I.18, I.35		145, 152, 167	
<i>Leptotila verreauxi</i>	I.38, I.42		90, 138, 167	
Psittacidae				
<i>Primolius maracana</i>		II.6	148	
<i>Diopsittaca nobilis</i>	I.27		167	
<i>Aratinga leucophthalma</i>	I.20, I.25, I.26, I.32, I.36, I.42, I.43, I.44, I.49	II.6, II.7, II.8	90, 139, 143, 148, 152, 167	
<i>Aratinga auricapillus</i>	I.27		167	
<i>Aratinga aurea</i>	I.24, I.27, I.32, I.35, I.38, I.41, I.46, I.47, I.50		47, 84, 138, 139, 145, 152, 159, 167	
<i>Aratinga cactorum</i>	I.3, I.5, I.10, I.15, I.19, I.20, I.21, I.23, I.24, I.25		70, 71, 82, 106, 139, 167	
<i>Pyrrhura frontalis</i>	I.5, I.15, I.27	II.8, III.2	5, 8, 16, 19, 71, 82, 136, 166, 167	AF
<i>Forpus xanthopterygius</i>	I.35		47, 84, 167	
<i>Brotogeris chiriri</i>	I.24, I.26, I.27, I.36		139, 167	
<i>Pionopsitta pileata</i>		II.8	7, 8, 11, 16, 19	AF
<i>Pionus maximiliani</i>	I.20, I.26, I.27, I.42, I.43		90, 139, 167	
<i>Amazona aestiva</i>	I.5		106	
Cuculidae				
<i>Piaya cayana</i>	I.5, I.27, I.35, I.40, I.42		71, 82, 90, 145, 164, 167	
<i>Crotophaga ani</i>	I.5, I.10, I.15, I.32, I.35, I.42, I.46		47, 71, 82, 84, 90, 106, 167	
<i>Guira guira</i>	I.26, I.35	II.1, II.8	10, 11, 16, 19, 84, 102, 139, 145, 167	
Tytonidae				
<i>Tyto alba</i>		II.6	148	
Strigidae				
<i>Megascops choliba</i>		II.6	148	
<i>Glaucidium brasilianum</i>	I.26, I.27, I.32		139, 167	
<i>Athene cunicularia</i>	I.18, I.33, I.35, I.46	II.6	47, 84, 148, 167	
Nyctibiidae				
<i>Nyctibius griseus</i>	I.35		84, 167	

Family/Species	<i>Campos rupestres</i>	<i>Campos de altitude</i>	Sources	Endemism
<b>Caprimulgidae</b>				
<i>Chordeiles pusillus</i>	I.25, I.43		139, 167	
<i>Caprimulgus longirostris</i>	I.10, I.20, I.26, I.27, I.30, I.32, I.35, I.36, I.38, I.40, I.42, I.44, I.46, I.49, I.50	II.1, II.2, II.6, II.8, II.10, III.2	20, 26, 28, 58, 67, 69, 70, 71, 76, 82, 84, 90, 102, 136, 138, 139, 145, 148, 152, 155, 156, 157, 159, 163, 164, 165, 167	
<i>Caprimulgus parvulus</i>	I.35		145	
<i>Hydropsalis torquata</i>	I.35, I.42, I.46, I.50	II.6	84, 90, 148, 152, 167	
<b>Apodidae</b>				
<i>Cypseloides senex</i>	I.15, I.35		70, 71, 82, 152, 167	
<i>Streptoprocne zonaris</i>	I.10, I.15, I.21, I.35, I.38, I.40, I.42, I.50	II.6, II.8, III.2	11, 16, 19, 70, 71, 82, 84, 90, 106, 136, 138, 145, 148, 152, 162, 164, 167	
<i>Streptoprocne biscutata</i>	I.5, I.10, I.20, I.21, I.26, I.27, I.32, I.35, I.36, I.38, I.42, I.43, I.46, I.51	II.6, II.8, III.2	5, 8, 70, 71, 82, 88, 90, 136, 148, 152, 156, 159, 165, 166, 167	
<i>Chaetura meridionalis</i>	I.38, I.40, I.42	II.6, II.12	90, 138, 148, 164, 167	
<b>Trochilidae</b>				
<i>Phaethornis pretrei</i>	I.3, I.5, I.8, I.10, I.15, I.18, I.20, I.21, I.23, I.24, I.25, I.26, I.27, I.30, I.32, I.33, I.35, I.36, I.38, I.42, I.43, I.44, I.46, I.47, I.48, I.50, II.5	II.1, II.3, II.13	6, 10, 25, 29, 33, 47, 50, 56, 68, 70, 71, 72, 75, 82, 84, 89, 90, 96, 98, 99, 100, 102, 106, 109, 126, 134, 135, 138, 139, 145, 152, 167	
<i>Phaethornis eurynome</i>		II.6, II.8	7, 16, 19, 148, 167	AF
<i>Campylopterus largipennis</i>	I.24, I.27, I.30, I.32, I.35, I.42		6, 25, 27, 50, 64, 89, 90, 139, 151, 152, 153, 167	
<i>Eupetomena macroura</i>	I.3, I.8, I.15, I.24, I.25, I.26, I.27, I.30, I.32, I.35, I.36, I.38, I.42, I.43, I.47, I.50	II.3, II.7	6, 25, 47, 50, 70, 71, 72, 74, 75, 82, 84, 89, 90, 91, 109, 118, 139, 145, 152, 167	
<i>Florisuga fusca</i>	I.30, I.38, I.42, I.45	II.3	25, 72, 75, 89, 90, 138, 158, 167	AF
<i>Colibri delphinae</i>	I.5, I.8, I.9, I.10		22, 50, 68, 70, 71, 82, 151	
<i>Colibri serrirostris</i>	I.3, I.5, I.9, I.10, I.15, I.18, I.20, I.23, I.24, I.25, I.26, I.27, I.30, I.32, I.33, I.35, I.36, I.38, I.42, I.43, I.44, I.45, I.46, I.47, I.48, I.50, I.51	II.1, II.2, II.3, II.6, II.7, II.8, II.11, II.13, III.1, III.2	6, 7, 8, 11, 16, 19, 25, 29, 34, 39, 44, 47, 50, 52, 56, 58, 68, 69, 70, 71, 72, 75, 80, 82, 83, 84, 89, 90, 96, 98, 102, 106, 126, 134, 136, 138, 139, 143, 145, 148, 152, 159, 162, 167	
<i>Anthracothorax nigricollis</i>		II.3	72	
<i>Chrysolampis mosquitus</i>	I.3, I.5, I.8, I.10, I.20, I.35		50, 68, 70, 71, 82, 96, 98, 100, 106, 134, 145, 152, 167	
<i>Stephanoxis lalandi</i>		II.1, II.2, II.3, II.6, II.8, II.10, II.11, II.12, III.2, III.3	5, 6, 7, 8, 10, 11, 16, 17, 19, 50, 52, 58, 69, 72, 102, 136, 137, 148, 151, 152, 159, 162, 166, 167	AF
<i>Chlorostilbon lucidus</i>	I.3, I.5, I.8, I.10, I.15, I.18, I.20, I.24, I.27, I.30, I.35, I.36, I.37, I.38, I.40, I.42, I.43, I.44, I.46, I.47, I.48, I.50	II.1, II.3, II.7, II.8, II.13	6, 10, 17, 25, 29, 44, 47, 50, 56, 68, 70, 71, 72, 74, 75, 82, 84, 86, 89, 90, 96, 98, 99, 100, 102, 106, 109, 126, 134, 138, 139, 143, 145, 152, 164, 167	
<i>Thalurania furcata</i>	I.30, I.35, I.42		25, 90, 167	
<i>Thalurania glaucopis</i>	I.8, I.42, I.50	II.11, II.13	50, 56, 89, 90, 167	AF
<i>Hylocharis cyanus</i>		II.1, II.3	17, 72	
<i>Leucochloris albicollis</i>	I.40, I.42, I.50	II.1, II.2, II.3, II.6, II.7, II.8, II.11, II.13, III.1, III.2	5, 6, 7, 8, 10, 11, 16, 17, 19, 52, 56, 72, 80, 90, 102, 136, 143, 148, 152, 162, 167	AF
<i>Amazilia versicolor</i>	I.8, I.30, I.36		25, 50, 167	
<i>Amazilia fimbriata</i>	I.3, I.8, I.24, I.27, I.30		25, 50, 68, 152, 167	

Family/Species	<i>Campos rupestres</i>	<i>Campos de altitude</i>	Sources	Endemism
<i>Amazilia lactea</i>	I.10, I.30, I.35, I.36, I.38, I.40, I.42, I.43, I.45, I.46, I.50	II.3	6, 18, 25, 72, 75, 84, 89, 90, 96, 100, 106, 134, 138, 145, 159, 164, 167	
<i>Clytolaema rubricauda</i>	I.42	II.1, II.3, II.7, II.8, II.11, II.12, III.2	5, 6, 7, 8, 10, 11, 16, 17, 19, 52, 72, 90, 102, 136, 143, 159, 167	AF
<i>Augastes scutatus</i>	I.20, I.24, I.25, I.26, I.27, I.29, I.30, I.32, I.33, I.35, I.36, I.37, I.39, I.40, I.42, I.43, I.44, I.45, I.46, I.47, I.48, I.50, I.51		6, 17, 18, 21, 23, 24, 25, 29, 30, 34, 37, 40, 42, 44, 47, 50, 58, 60, 61, 62, 64, 74, 77, 81, 84, 89, 90, 91, 93, 95, 107, 108, 115, 129, 131, 139, 145, 151, 152, 153, 157, 158, 159, 160, 161, 164, 167	MT
<i>Augastes lumachella</i>	I.1, I.2, I.3, I.4, I.5, I.8, I.9, I.10, I.11, I.12, I.13, I.14, I.15, I.16, I.17		21, 23, 24, 29, 30, 38, 50, 58, 68, 70, 71, 82, 85, 96, 100, 101, 106, 107, 134, 151, 153, 159, 167	MT
<i>Heliactin bilophus</i>	I.9, I.15, I.27, I.30, I.33, I.35		25, 68, 70, 71, 82, 84, 89, 145, 159, 167	
<i>Calliphlox amethystina</i>	I.8, I.10, I.30, I.35, I.36, I.42	II.3	25, 29, 47, 50, 72, 89, 90, 96, 98, 100, 106, 134, 167	
Bucconidae				
<i>Nystalus chacuru</i>	I.15, I.32, I.35, I.46, I.48, I.49		84, 145, 152, 167	
Ramphastidae				
<i>Ramphastos toco</i>		II.6	148	
Picidae				
<i>Picumnis cirratus</i>	I.35, I.36, I.40, I.42, I.50		90, 164, 167	
<i>Melanerpes candidus</i>	I.32, I.35	II.1, II.6	10, 47, 102, 148, 167	
<i>Veniliornis mixtus</i>	I.35		145	
<i>Colaptes campestris</i>	I.5, I.10, I.15, I.21, I.23, I.24, I.25, I.26, I.27, I.32, I.33, I.35, I.36, I.37, I.38, I.40, I.42, I.43, I.44, I.46, I.47, I.48, I.50, II.5	II.1, II.6, II.7, II.8	5, 7, 8, 10, 11, 16, 19, 39, 47, 71, 82, 84, 90, 102, 106, 138, 139, 143, 145, 148, 164, 167	
Melanopareiidae				
<i>Melanopareia torquata</i>	I.5, I.10, I.15, I.19, I.20, I.25, I.27, I.34, I.35, I.38, I.44, I.46, I.49, I.50		53, 62, 70, 71, 77, 82, 84, 139, 152, 167	CE
Thamnophilidae				
<i>Mackenziaena leachii</i>	I.35, I.36, I.42, I.46, I.48	II.1, II.8, III.2	7, 8, 11, 14, 16, 18, 19, 58, 60, 78, 90, 102, 136, 159, 162, 166, 167	AF
<i>Tanaba major</i>	I.27		167	
<i>Sakesphorus cristatus</i>	I.3, I.20, I.24, I.26		70, 71, 139, 167	CA
<i>Thamnophilus doliatus</i>	I.23, I.24		139, 167	
<i>Thamnophilus caerulescens</i>		II.6, II.8, II.10, III.2	5, 7, 11, 16, 19, 136, 137, 148, 165, 166	
<i>Thamnophilus torquatus</i>	I.5, I.10, I.15, I.19, I.20, I.35, I.36		70, 71, 77, 82, 84, 106, 145, 167	
<i>Thamnophilus ruficapillus</i>	I.42, I.48, I.50	II.1, II.6, II.8, II.10, III.2	10, 11, 14, 16, 18, 19, 43, 58, 65, 69, 90, 102, 136, 137, 148, 162, 166, 167	
<i>Myrmorchilus strigilatus</i>	I.15, I.20, I.21, I.24, I.26		139, 152, 167	
<i>Herpsilochmus atricapillus</i>	I.20, I.27		167	
<i>Formicivora serrana</i>	I.36, I.42, I.43		77, 90, 167	AF
<i>Formicivora melanogaster</i>	I.20		167	
<i>Formicivora rufa</i>	I.15, I.26, I.27		70, 71, 82, 139, 167	
<i>Formicivora grantsaui</i>	I.5, I.6, I.7, I.9, I.10		70, 71, 82, 132	MT
<i>Drymophila genei</i>		II.1, II.2, II.8, II.10, III.1, III.2	7, 11, 15, 16, 18, 19, 58, 65, 102, 136, 137, 159, 162, 166, 167	AF



Family/Species	<i>Campos rupestres</i>	<i>Campos de altitude</i>	Sources	Endemism
Conopophagidae				
<i>Conopophaga lineata</i>		II.1, II.8, III.2	5, 7, 8, 11, 14, 16, 18, 19, 28, 52, 58, 102, 136, 151, 159, 165, 166	
Rhinocryptidae				
<i>Scytalopus notorius</i>		II.1, II.2, II.6, II.7, II.8, II.10, II.11, II.12, III.2	5, 7, 8, 11, 12, 13, 16, 18, 19, 28, 39, 45, 52, 58, 102, 124, 136, 137, 148, 151, 152, 157, 159, 162, 166, 167	AF
<i>Scytalopus speluncae</i>	I.32, I.35, I.36, I.37, I.40, I.42, I.44, I.46, I.48		47, 77, 84, 90, 152, 155, 159, 167	
<i>Scytalopus iraiensis</i>	I.35, I.42		149, 167	AF
Dendrocolaptidae				
<i>Lepidocolaptes angustirostris</i>	I.20, I.21, I.24, I.25, I.26, I.27, I.35, I.42		47, 84, 90, 139, 167	
Furnariidae				
<i>Cinclodes pabsti</i>	I.35		147, 152	AF
<i>Furnarius rufus</i>	I.15, I.26, I.27, I.32, I.35, I.36, I.40, I.46, I.48	II.1	10, 47, 84, 102, 139, 164, 167	
<i>Oreophylax moreirae</i>	I.42	II.1, II.8, II.9, II.10, III.2	5, 7, 8, 9, 10, 11, 16, 19, 32, 39, 43, 52, 58, 64, 69, 79, 81, 90, 102, 104, 136, 137, 141, 151, 152, 155, 156, 157, 159, 162, 165, 166, 167	MT
<i>Synallaxis frontalis</i>	I.3, I.5, I.15, I.18, I.20, I.24, I.26, I.27, I.35		71, 82, 84, 139, 152, 167	
<i>Synallaxis albescens</i>	I.23, I.24, I.25, I.26, I.35, I.50		84, 139, 145, 167	
<i>Synallaxis spixi</i>	I.5, I.10, I.15, I.20, I.21, I.25, I.26, I.27, I.32, I.35, I.36, I.38, I.40, I.42, I.44, I.46, I.48, I.50	II.1, II.3, II.6, II.7, II.8, II.11, III.2	39, 47, 70, 71, 72, 82, 84, 90, 102, 104, 136, 138, 139, 148, 152, 162, 164, 166, 167	AF
<i>Cranioleuca pallida</i>	I.42	II.1, II.8, III.2	5, 8, 11, 16, 19, 90, 136, 166, 167	
<i>Asthenes luizae</i>	I.21, I.25, I.26, I.32, I.33, I.34, I.35		42, 46, 48, 61, 62, 84, 93, 94, 114, 115, 116, 117, 119, 139, 145, 150, 152, 156, 157, 159, 167	MT
<i>Phacellodomus rufifrons</i>	I.15, I.21, I.25, I.26, I.33, I.35, I.46, I.50	II.1	47, 84, 102, 139, 167	
<i>Anumbius annumbi</i>	I.35, I.37, I.48, I.51	II.7	84, 167	
<i>Lochmias nematura</i>	I.21, I.23, I.25, I.32, I.35, I.42, I.46, I.48	II.1, II.6, II.7, II.8, III.2	7, 8, 10, 11, 16, 19, 84, 102, 136, 148, 152, 159, 166, 167	
Tyrannidae				
<i>Hemitriccus obsoletus</i>		II.8, II.10	5, 7, 8, 11, 16, 19, 137, 159	AF
<i>Hemitriccus margaritaceiventer</i>	I.10, I.15, I.18, I.20, I.25, I.26, I.27, I.32, I.35		71, 82, 106, 139, 145, 152, 167	
<i>Todirostrum cinereum</i>	I.15		167	
<i>Phyllomyias fasciatus</i>	I.32, I.35, I.36, I.42, I.43, I.44, I.46	II.8	7, 11, 16, 19, 47, 84, 152, 159, 167	
<i>Elaenia flavogaster</i>	I.15, I.20, I.21, I.24, I.25, I.26, I.27, I.32, I.33, I.35, I.38, I.40, I.42, I.43, I.47, I.48, I.50		47, 70, 71, 84, 90, 138, 139, 145, 152, 164, 167	
<i>Elaenia albiceps</i>		II.6	148	
<i>Elaenia parvirostris</i>	I.43		167	
<i>Elaenia mesoleuca</i>	I.42, I.44	II.8, III.2	11, 16, 19, 43, 136, 152, 159, 162, 167	
<i>Elaenia cristata</i>	I.5, I.10, I.15, I.18, I.20, I.21, I.23, I.24, I.25, I.26, I.27, I.32, I.33, I.35, I.50, II.5		47, 70, 71, 82, 84, 113, 118, 139, 145, 152, 167	
<i>Elaenia chiriquensis</i>	I.20, I.27, I.35, I.42, I.43, I.50		84, 90, 145, 167	

Family/Species	<i>Campos rupestres</i>	<i>Campos de altitude</i>	Sources	Endemism
<i>Elaenia obscura</i>	I.15, I.24, I.25, I.26, I.27, I.32, I.35, I.36, I.40, I.42, I.43, I.44, I.46, I.49, I.50	II.6, II.8, II.10, II.11, III.2	5, 8, 16, 19, 47, 84, 90, 104, 136, 137, 139, 145, 148, 152, 159, 164, 165, 166, 167	
<i>Camptostoma obsoletum</i>	I.3, I.5, I.10, I.15, I.18, I.20, I.21, I.23, I.24, I.25, I.26, I.27, I.35, I.36, I.38, I.40, I.42, I.43, I.46, I.50	II.1, II.6, II.7	47, 71, 82, 84, 90, 138, 139, 143, 145, 148, 152, 164, 167	
<i>Serpophaga nigricans</i>	I.42	II.8	86, 159, 167	
<i>Serpophaga subcristata</i>	I.35, I.42, I.43	II.1, II.7, II.8	7, 11, 16, 19, 90, 102, 167	
<i>Phaeomyias murina</i>	I.20, I.24, I.32, I.35, I.42, I.43		84, 90, 139, 167	
<i>Polystictus superciliosus</i>	I.3, I.5, I.15, I.20, I.22, I.25, I.26, I.32, I.33, I.35, I.36, I.37, I.38, I.39, I.40, I.42, I.43, I.44, I.46, I.48, I.50	II.6, II.7, II.8, III.3	2, 35, 38, 42, 47, 49, 53, 58, 60, 62, 66, 70, 71, 73, 74, 77, 81, 82, 84, 90, 93, 104, 105, 115, 119, 120, 121, 122, 123, 133, 138, 139, 143, 145, 148, 151, 152, 159, 164, 167	MT
<i>Phylloscartes ventralis</i>	I.40	II.6, II.7, II.8, II.11	5, 7, 8, 11, 16, 19, 43, 148, 164, 167	
<i>Phylloscartes difficilis</i>		II.8, III.2	5, 7, 8, 11, 16, 19, 52, 136, 165, 166	AF
<i>Culicivora caudacuta</i>	I.35		167	
<i>Myiophobus fasciatus</i>	I.18, I.20, I.25, I.26, I.27, I.32, I.35, I.36, I.38, I.40, I.42, I.43, I.46, I.48, I.50	II.1, II.7	10, 47, 84, 90, 102, 138, 139, 143, 145, 152, 167	
<i>Hirundinea ferruginea</i>	I.3, I.5, I.10, I.15, I.20, I.23, I.24, I.25, I.27, I.32, I.35, I.36, I.38, I.42, I.43, I.44, I.46, I.48, I.50, II.5	II.2, II.3, II.6, II.11	60, 70, 71, 72, 82, 84, 90, 106, 138, 139, 148, 152, 167	
<i>Pyrocephalus rubinus</i>		II.3	72	
<i>Knipolegus cyanirostris</i>	I.35, I.42	II.1, II.2, II.6, II.8, II.10	5, 7, 8, 10, 11, 16, 19, 52, 90, 102, 137, 148, 167	
<i>Knipolegus lophotes</i>	I.26, I.32, I.35, I.36, I.38, I.40, I.42, I.44, I.45, I.48, I.50, I.51, II.5	II.1, II.6, II.7, II.8	7, 8, 11, 16, 18, 19, 47, 58, 60, 84, 90, 102, 138, 139, 143, 145, 148, 159, 164, 167	
<i>Knipolegus nigerrimus</i>	I.3, I.5, I.10, I.15, I.20, I.21, I.23, I.24, I.25, I.26, I.27, I.29, I.32, I.33, I.35, I.36, I.37, I.38, I.40, I.41, I.42, I.43, I.44, I.46, I.48, I.49, I.50, I.51, II.5	II.1, II.2, II.6, II.7, II.8, II.10, II.11, II.12, III.1, III.2	4, 5, 7, 8, 10, 11, 16, 19, 51, 54, 58, 66, 69, 70, 71, 82, 84, 90, 92, 97, 102, 104, 106, 113, 119, 136, 137, 138, 139, 143, 145, 148, 151, 152, 159, 162, 164, 165, 166, 167	
<i>Xolmis cinereus</i>	I.10, I.24, I.25, I.26, I.27, I.32, I.35, I.38, I.42, I.46	II.8	4, 5, 8, 11, 16, 19, 47, 70, 71, 82, 84, 90, 138, 139, 145, 167	
<i>Xolmis velatus</i>	I.25, I.35, I.40, I.43, I.51	II.6, II.7, II.8	7, 8, 47, 84, 143, 145, 148, 152, 164, 167	
<i>Muscipira vetula</i>	I.35, I.42, I.46, I.48	II.1, II.8, II.11	5, 8, 10, 11, 16, 19, 60, 90, 102, 167	AF
<i>Myiozetetes similis</i>	I.15, I.26, I.42		90, 139, 167	
<i>Pitangus sulphuratus</i>	I.5, I.10, I.24, I.32, I.35, I.38, I.40, I.42, I.46	II.1, II.8	7, 10, 11, 16, 19, 47, 84, 90, 102, 106, 138, 139, 145, 164, 167	
<i>Megarynchus pitangua</i>	I.5, I.15, I.40, I.42		90, 106, 164, 167	
<i>Tyrannus albogularis</i>	I.40		164	
<i>Tyrannus melancholicus</i>	I.5, I.10, I.15, I.24, I.32, I.35, I.38, I.40, I.42, I.43, I.50	II.6, II.8	5, 8, 11, 16, 19, 71, 84, 90, 106, 145, 148, 152, 164, 167	
<i>Tyrannus savana</i>	I.35, I.38, I.40, I.43		138, 145, 164, 167	
<i>Myiarchus swainsoni</i>	I.27, I.42		90, 167	
<i>Myiarchus ferrox</i>	I.26, I.27, I.35, I.36, I.38, I.42, I.46, I.48, I.50	II.6	47, 84, 90, 139, 148, 152, 167	
<i>Myiarchus tyrannulus</i>	I.42		90	

Family/Species	<i>Campos rupestres</i>	<i>Campos de altitude</i>	Sources	Endemism
<i>Ramphotrigon megalcephalum</i>		II.1	18, 102	
Tityridae				
<i>Tityra cayana</i>	I.27		167	
Vireonidae				
<i>Cyclarhis gujanensis</i>	I.10, I.18, I.20, I.21, I.23, I.24, I.25, I.27, I.32, I.35, I.36, I.38, I.40, I.42, I.43, I.46, I.50	II.1, II.6, II.7, II.8, III.2	5, 7, 10, 11, 16, 19, 28, 47, 52, 84, 90, 97, 102, 106, 113, 136, 138, 145, 148, 164, 166, 167	
<i>Hylophilus amaurocephalus</i>	I.21		167	
Corvidae				
<i>Cyanocorax cristatellus</i>	I.25, I.27, I.32, I.35, I.38	II.6	84, 138, 145, 148, 167	CE
<i>Cyanocorax cyanopogon</i>	I.27		167	
Hirundinidae				
<i>Progne tapera</i>	I.5, I.10, I.24	II.6	106, 148, 167	
<i>Progne chalybea</i>	I.24, I.36, I.38, I.51		138, 167	
<i>Pygochelidon cyanoleuca</i>	I.5, I.10, I.15, I.20, I.21, I.24, I.26, I.35, I.36, I.37, I.38, I.42, I.44, I.46, I.47, I.50, I.51	II.1, II.6, II.7, II.8, II.11, II.12, III.1, III.2	5, 11, 16, 19, 47, 58, 70, 71, 82, 84, 90, 136, 138, 139, 143, 145, 148, 152, 165, 167	
<i>Alopochelidon fucata</i>	I.25, I.35, I.38		47, 84, 138, 167	
<i>Stelgidopteryx ruficollis</i>	I.15, I.21, I.24, I.25, I.26, I.27, I.32, I.35, I.36, I.38, I.40, I.42, I.43, I.46, I.50		47, 71, 82, 84, 90, 138, 139, 145, 164, 167	
Troglodytidae				
<i>Troglodytes musculus</i>	I.5, I.10, I.15, I.18, I.20, I.21, I.23, I.24, I.25, I.26, I.27, I.32, I.33, I.35, I.36, I.38, I.40, I.42, I.43, I.44, I.46, I.48, I.50, II.5	II.6, II.7, II.11	47, 71, 82, 84, 90, 106, 138, 139, 145, 148, 152, 164, 167	
<i>Pheugopedius genibarbis</i>	I.5, I.10, I.15		70, 71, 82, 167	
Poliptilidae				
<i>Poliptila plumbea</i>	I.3, I.10, I.15, I.18, I.20, I.25		70, 71, 139, 167	
Turdidae				
<i>Turdus flavipes</i>	I.35, I.42	II.8	5, 7, 8, 11, 16, 19, 43, 52, 145, 167	
<i>Turdus rufiventris</i>	I.5, I.27, I.35, I.40, I.42	II.1, II.6, II.8, III.2	7, 10, 11, 16, 19, 52, 84, 102, 106, 136, 148, 159, 164, 166, 167	
<i>Turdus leucomelas</i>	I.3, I.5, I.10, I.15, I.18, I.20, I.21, I.24, I.25, I.26, I.27, I.32, I.35, I.36, I.38, I.42, I.46, I.50, II.5	II.6, II.8	7, 11, 47, 71, 82, 84, 90, 97, 106, 113, 138, 139, 145, 148, 152, 167	
<i>Turdus amaurochalinus</i>	I.27, I.35, I.40	II.7, II.8	16, 19, 84, 159, 164, 167	
<i>Turdus albicollis</i>		II.6	148	
Mimidae				
<i>Mimus saturninus</i>	I.5, I.10, I.15, I.21, I.23, I.24, I.26, I.27, I.32, I.35, I.36, I.37, I.38, I.40, I.42, I.46, I.43, I.47, I.48, I.50	II.1, II.6, II.7	10, 47, 70, 71, 84, 90, 102, 106, 145, 148, 164, 167	
Motacillidae				
<i>Anthus hellmayri</i>	I.32, I.35, I.46, I.48, I.49, I.50, I.51	II.1, II.6, II.7, II.8, III.2	5, 7, 8, 11, 16, 19, 39, 47, 54, 55, 58, 84, 102, 125, 136, 143, 145, 148, 152, 159, 165, 167	
Coerebidae				
<i>Coereba flaveola</i>	I.3, I.5, I.10, I.15, I.18, I.20, I.24, I.25, I.26, I.27, I.35, I.36, I.38, I.42, I.43, I.48, I.50	II.1	10, 28, 71, 82, 84, 90, 98, 102, 106, 113, 126, 138, 139, 152, 167	

Family/Species	<i>Campos rupestres</i>	<i>Campos de altitude</i>	Sources	Endemism
Thraupidae				
<i>Schistochlamys ruficapillus</i>	I.3, I.5, I.10, I.15, I.18, I.20, I.21, I.24, I.25, I.26, I.27, I.29, I.32, I.33, I.35, I.36, I.38, I.37, I.40, I.41, I.42, I.43, I.44, I.46, I.48, I.50	II.1, II.3, II.6, II.8, III.2	10, 47, 52, 70, 71, 72, 82, 84, 90, 97, 102, 106, 112, 113, 118, 119, 136, 138, 139, 145, 148, 152, 159, 162, 164, 167	
<i>Cypsnagra hirundinacea</i>	I.26, I.35		84, 115, 139, 145, 167	
<i>Trichothraupis melanops</i>	I.50		152, 167	
<i>Piranga flava</i>	I.5, I.10, I.15, I.24, I.27, I.32, I.35, I.36, I.38, I.42, I.50	II.6, II.8	5, 11, 16, 19, 70, 71, 82, 84, 90, 97, 106, 113, 138, 145, 148, 167	
<i>Tachyphonus rufus</i>	I.20		167	
<i>Thraupis sayaca</i>	I.5, I.10, I.20, I.24, I.26, I.27, I.35, I.36, I.38, I.40, I.42, I.43, II.5		47, 84, 90, 106, 139, 164, 167	
<i>Stephanophorus diadematus</i>		II.1, II.2, II.6, II.7, II.8, II.10, II.11, II.12, III.1, III.2	5, 7, 8, 10, 11, 16, 18, 19, 43, 52, 58, 69, 73, 102, 104, 136, 137, 143, 148, 152, 159, 162, 166, 167	AF
<i>Pipraeidea melanonota</i>	I.35, I.42, I.50	II.8	5, 8, 11, 16, 19, 90, 145, 152, 167	
<i>Tangara desmaresti</i>	I.40, I.42	II.1, II.6, II.7, II.8, II.11, III.2	7, 8, 10, 11, 16, 19, 43, 90, 102, 136, 143, 148, 152, 159, 164, 166, 167	AF
<i>Tangara cyanoventris</i>	I.10, I.15, I.36, I.38, I.40, I.42, I.49		66, 71, 77, 82, 90, 138, 164, 152, 167	AF
<i>Tangara cayana</i>	I.5, I.10, I.15, I.18, I.20, I.21, I.24, I.26, I.27, I.32, I.35, I.36, I.38, I.42, I.43, I.46, I.48, I.50, II.5	II.6, II.7, II.11	47, 71, 82, 84, 90, 106, 138, 139, 143, 145, 148, 152, 167	
<i>Tersina viridis</i>	I.35, I.42		90, 118	
<i>Dacnis cayana</i>	I.25, I.27, I.36, I.38, I.40, I.42		90, 138, 139, 164, 167	
<i>Hemithraupis guira</i>	I.27		167	
<i>Conirostrum speciosum</i>	I.48		167	
Emberizidae				
<i>Zonotrichia capensis</i>	I.3, I.5, I.10, I.15, I.18, I.20, I.21, I.23, I.24, I.25, I.26, I.27, I.32, I.33, I.35, I.36, I.37, I.38, I.40, I.42, I.43, I.44, I.46, I.47, I.48, I.50, I.51, II.5	II.1, II.6, II.7, II.8, II.10, II.11, II.12, III.2, III.3	4, 5, 7, 8, 10, 11, 16, 19, 39, 43, 47, 52, 58, 68, 71, 82, 84, 90, 97, 102, 106, 113, 136, 137, 138, 139, 143, 145, 148, 151, 152, 159, 162, 164, 166, 167	
<i>Ammodramus humeralis</i>	I.5, I.10, I.15, I.20, I.24, I.25, I.26, I.27, I.32, I.35, I.40, I.43, I.48	II.6	47, 70, 71, 82, 84, 139, 145, 148, 152, 164, 167	
<i>Porphyrospiza caerulescens</i>	I.15, I.20, I.24, I.25, I.27, I.32, I.35, I.38, I.39, I.40, I.47		53, 70, 71, 77, 82, 84, 115, 139, 145, 152, 164, 167	CE
<i>Haplospiza unicolor</i>	I.36, I.42	II.1, II.8	10, 11, 16, 19, 43, 90, 102, 152, 167	AF
<i>Donacospiza albifrons</i>	I.35, I.42, I.48	II.1, II.8, III.2	16, 19, 58, 102, 136, 159, 166, 167	
<i>Poospiza thoracica</i>		II.1, II.7, II.8, II.10, II.11, II.12, III.2, III.3	4, 5, 7, 8, 10, 11, 16, 19, 52, 73, 102, 136, 137, 143, 152, 158, 159, 162, 166, 167	AF
<i>Poospiza lateralis</i>		II.1, II.7, II.8, II.11, II.12, III.3	7, 8, 10, 11, 16, 19, 52, 69, 102, 130, 143, 151, 159, 157, 167	AF
<i>Poospiza cinerea</i>	I.27, I.35, I.38		53, 111, 115, 119, 145, 167	CE
<i>Sicalis citrina</i>	I.5, I.10, I.15, I.18, I.20, I.21, I.23, I.24, I.25, I.26, I.27, I.32, I.34, I.35, I.36, I.37, I.38, I.42, I.43, I.46, I.48, I.49, I.50, II.5	II.2, II.6, II.7, II.11, II.12	60, 62, 70, 71, 82, 84, 90, 138, 139, 140, 143, 144, 145, 148, 152, 167	

Family/Species	<i>Campos rupestres</i>	<i>Campos de altitude</i>	Sources	Endemism
<i>Sicalis flaveola</i>	I.5, I.10	II.1, II.6, II.8	7, 8, 10, 11, 16, 19, 54, 55, 102, 106, 167	
<i>Emberizoides herbicola</i>	I.15, I.24, I.25, I.26, I.27, I.35, I.38, I.42, I.46, I.48, I.50, II.5	II.6, II.8	7, 8, 11, 16, 19, 47, 54, 55, 70, 71, 82, 84, 86, 104, 139, 145, 148, 167	
<i>Emberizoides ypiranganus</i>	I.32, I.35		127, 146, 152, 167	AF
<i>Embernagra platensis</i>		II.1, II.8, II.10	5, 8, 11, 16, 19, 38, 43, 58, 102, 137, 167	
<i>Embernagra longicauda</i>	I.3, I.5, I.10, I.15, I.19, I.20, I.21, I.23, I.24, I.25, I.26, I.27, I.28, I.29, I.30, I.31, I.32, I.33, I.34, I.35, I.36, I.38, I.39, I.40, I.42, I.43, I.44, I.45, I.46, I.47, I.48, I.50, I.51, II.4, II.5	II.1	31, 35, 36, 38, 41, 42, 47, 53, 58, 60, 61, 62, 63, 64, 70, 71, 74, 77, 81, 82, 84, 87, 90, 93, 102, 103, 104, 115, 119, 127, 138, 139, 142, 145, 151, 152, 154, 157, 159, 167	MT
<i>Volatinia jacarina</i>	I.5, I.10, I.24, I.35, I.38, I.40, I.43, I.46, I.50		70, 71, 82, 84, 106, 138, 139, 145, 152, 164, 167	
<i>Sporophila plumbea</i>	I.24, I.25		139, 167	
<i>Sporophila nigricollis</i>	I.5, I.10, I.20, I.24, I.25, I.27, I.32, I.35, I.36, I.40, I.42, I.43		84, 90, 106, 139, 164, 167	
<i>Sporophila caeruleascens</i>	I.35	II.1	47, 84, 102, 167	
<i>Sporophila bouvreuil</i>	I.25		167	
<i>Coryphospingus pileatus</i>	I.35, I.43, I.50	II.1	10, 84, 102, 167	
Cardinalidae				
<i>Saltator similis</i>	I.10, I.15, I.20, I.21, I.27, I.35, I.38, I.40, I.42		90, 97, 113, 138, 164, 167	
<i>Saltator maxillosus</i>		II.8, III.3	5, 7, 8, 11, 16, 19, 43, 159	AF
<i>Saltator atricollis</i>	I.3, I.5, I.10, I.15, I.26, I.27, I.32, I.35, I.42		47, 68, 70, 71, 82, 84, 90, 115, 139, 145, 152, 167	CE
Parulidae				
<i>Parula pitiayumi</i>	I.20, I.27		167	
<i>Geothlypis aequinoctialis</i>	I.5, I.10, I.15, I.25, I.26, I.27, I.35, I.38, I.42, I.43, I.44, I.46, I.48, I.50	II.1	28, 69, 70, 71, 82, 84, 90, 102, 138, 139, 145, 152, 167	
Icteridae				
<i>Gnorimopsar chopi</i>	I.5, I.10, I.15, I.24, I.25, I.26, I.27, I.32, I.33, I.35, I.46	II.6	47, 70, 71, 82, 84, 106, 139, 145, 148, 152, 167	
<i>Pseudoleistes guinahuero</i>	I.35		84, 119, 145, 152, 167	
<i>Molothrus bonariensis</i>	I.26, I.35, I.36, I.37, I.38	II.8	7, 11, 16, 19, 46, 84, 110, 139, 145, 167	
Fringillidae				
<i>Carduelis magellanica</i>	I.5, I.24, I.35, I.50	II.6, II.7, II.8	5, 7, 8, 11, 16, 19, 43, 71, 82, 84, 139, 145, 148, 167	
<i>Euphonia chlorotica</i>	I.5, I.10, I.15, I.18, I.21, I.27, I.35, I.38, I.40, I.42, I.44		71, 82, 84, 90, 106, 164, 167	
<i>Euphonia cyanocephala</i>	I.24, I.35, I.42		139, 152, 167	
<i>Chlorophonia cyanea</i>	I.36, I.42, I.44		90, 152, 167	